



GREEN SYNTHESIS OF AN AMIDE-BASED CHEMOSENSOR AND ITS APPLICATION FOR DETECTION OF TOXIC METAL IONS

D. S. Bhagat,^{[a]*} R. P. Pawar,^{[b]*} A. B. Tekale,^[c] S. G. Pande,^[a] R. R. Rangari,^[a] I. V. Suryawanshi,^[a] P. B. Chavan^[a], S. N. Tapase^[a] and A. A. Sahu^[d]

Keywords: chemosensor, forensic application, ultra-sonication, coordination complex

Ultrasound-assisted synthesis of a chemosensor from benzoyl chloride and diethylenetriamine in acetonitrile was carried out and the product structure was confirmed by UV-VIS and FT-IR spectroscopy. The synthesized chemosensor interacts with metal ions and produces a notable change in colorimetric measurements. It is observed that as the concentration of metallic ions increases, absorbance also increases, thus the chemosensor can be used for quantitative estimation of metal-ion content.

* Corresponding Authors

E-Mail: rppawar@yahoo.com

- [a] Department of Forensic Chemistry and Toxicology, Government Institute of Forensic Science, Aurangabad-431005 (M.S.)
 [b] Department of Chemistry Deogiri College, Station Road, Aurangabad 431 005, Maharashtra, India
 [c] Department of Chemistry, Shri Shivaji College, Parbhani – 431 401 (MS) India.
 [d] Department of Forensic Chemistry and Toxicology, Government Institute of Forensic Science, Aurangabad-431005 (M.S.)

Introduction

In recent decade increase in industrialization resulted in high pollution of metal ions in the environment which interact with living systems and cause hazardous effects on human health and the environment resulting in toxic cation and poisoning.

Chemosensor is a sensory receptor that shows a detectable change in color and photophysical properties. The ultimate aim of the scientific community is the synthesis of sensitive chemosensor which can be used for the detection of heavy metal ions.¹ The Hg²⁺ is a common, environmentally toxic and hazardous pollutant among the heavy metal ions. Trace amounts of Hg²⁺ in the body can lead neurotoxicity and the mercury ions easily pass through the biological tissues to cause digestive, kidney and especially neurological diseases.² The Cu²⁺ ion is an important transitional metal ion which plays a major role in environmental, biological, forensic and chemical fields.³ Cu²⁺ ion is the third most abundant essential metal ion in the human body transition element after Fe³⁺ and Zn²⁺.⁴ Zinc is the second most abundant transition metal in the human body next to iron and acts as a catalytic co-factor in various metallic-enzymes. Zn²⁺ ions play a vital role in numerous fundamental biological processes such as cellular metabolism, DNA synthesis and neurotransmission.⁵ The deficiency of Zn²⁺ ions causes various diseases such as Alzheimer's, Parkinson's diseases, epilepsy regulation disorder in mammalian reproduction, immune system, sense of taste and smell and overall growth of the living body.⁶ The zinc is an important constituent in more than 250 metalloenzymes.⁷ Cobalt plays a significant role in the metabolism of ferrous


and hemoglobin synthesis. The Co²⁺ ions are an important constituent in vitamins.⁸ The Co²⁺ shows toxicological effects on human health, including decreased cardiac output, asthma, heart and lung disease.⁹ Ni²⁺ is an essential trace ion in organic systems, but the excess accumulation of Ni²⁺ in the body can lead to various diseases such as lung fibrosis, cardiovascular and kidney diseases.¹⁰

In recent decades, a wide number of efficient chemosensors (ligands) have been developed for the detection of various metal ions in aqueous solution. The type of interaction between chelate and the metal ion is a coordinate bond or host-guest type.¹¹ Various methods have been used for detection of the metal ions such as rhodamine 6G (R6G) and 8-aminoquinoline (8-AQ) co-modified core/shell Fe₃O₄@SiO₂ nanoparticles used for detection of Hg²⁺ and Zn²⁺ ions in aqueous solution,¹² Rhodamine B chemosensor used for detection of Pb²⁺ in acetonitrile,¹³ azadiene-pyrene derivative for Hg²⁺ in aqueous acetonitrile solution,¹⁴ (E)-2-(5-((2-carbamothioylhydrazono)methyl)-6-hydroxy-3-oxo-3H-xanthen-9-yl)benzoic acid for the detection of Cd²⁺ ions in aqueous solution,¹⁵ N-(1,10-phenanthroline-5-yl)-2,2-diphenylacetamide for selective detection of Fe³⁺ in an aqueous medium,¹⁶ 8-hydroxyquinoline based graphene oxide for detection of Zn²⁺ in aqueous media,¹⁷ cyclometalated platinum(II) bipyridylacetylidene complex for Mg²⁺ in acetonitrile,¹⁸ rhodamine-based compound for Pd²⁺ in pure water,¹⁹ and for highly selective detection of Ag(I) ions.²⁰

The syntheses of chemosensors are reported in the literature by various methods from readily available starting materials such as water mediated synthesis of CF₁-CF₃ and use for detection of CN⁻.²¹ Ultrasound-assisted synthesis of diphenylamine-based Schiff base used in the detection of copper(II) ion in aqueous solution,²² microwave-assisted synthesis of graphitic carbon nitride quantum dots²³ and photoluminescent carbon dots synthesis using starch (*Tapioca Sago*) via solution method at mild condition²⁴ and possess great applications in various fields such as environmental,²⁵ forensic,²⁶ and the biological sciences.²⁷

Herein, we develop an ultrasound-assisted new method for the synthesis of amide-based chemosensor using

diethylenediamine and benzoyl chloride. Ultrasound waves accelerate the rate of reaction by cavitation and nebulization phenomenon. The cavitation helps to create a development of implosive collapse of bubbles in a liquid. Nebulization phenomenon helps in the creation of mist from ultrasound passing through a liquid and impinging on a liquid-gas interface.²⁸



Materials and methods

All chemicals were purchased from Sigma Aldrich and used without further purification. UV spectrophotometric analysis was carried out using UV-VIS Perkin Elmer, lambda scan-35 system. FT-IR analysis was done on a BRUKER instrument having OPUS software of version 7.0.129. □

Procedure for synthesis of amide-based chemo-sensor

In 25 mL round bottom conical flask benzoyl chloride (5.00 mmol), diethylenetriamine (10.00 mmol) and 1-2 mL acetonitrile were taken and kept in ultrasonication bath at room temperature for 10 minutes. The reaction mixture was poured into ice-cold water, and the precipitate of chemosensor was separated by simple filtration. Residue allowed drying at room temperature and dry white crystalline powder compound is used as chemosensor in aqueous ethyl alcohol.

Procedure for preparation of pH 10 buffer solution

For 10 ml: pH 10 buffer was prepared by taking 0.7 g of NH₄Cl, and 5.68 mL NH₃ solution in 10 mL distilled water in a conical flask.

Results and discussion

In the present work, we developed a new method for the synthesis of amide-based chemosensor ((*7E*)-*N*¹-benzylidene-*N*²-((*E*)-2-(benzylideneamino)ethyl)ethane-1,2-diamine) using diethylenetriamine and benzoyl chloride under ultrasound irradiation. The absorbance in the UV-VIS analysis of Co(II) ion solution in aqueous ethanol for 100, 200, 300, 400, 500, 600, 700, 800, 900 and 1000 ppm are illustrated in Figure 2. It was observed that as the concentration of a metal ion in solution was increased, the absorbance was increased. It helps to find out the concentration of a metal ion in an unknown solution by

using a standard calibration curve. Mercury (Hg²⁺), nickel (Ni²⁺) and copper (Cu²⁺) show excellent absorbance in the UV-VIS analysis. It was observed that metal ions form stable complexes with chemosensor and show visible color change or change in color intensity of aqueous solution. In UV-VIS analysis, absorbance was increased with respect to concentration and it validates the Lambert-Beer's Law.

The FT-IR spectroscopic technique was used to predict the functional group of chemosensor and metal-complex of chemosensor with the help of stretching frequencies. These results are illustrated in Table 1. The probable interaction of chemosensor with metal ions shows that it forms a stable coordination complex with a metal ion (a predicted structure is given in Figure 1). The synthesized amide-based derivative probably acts as a tridentate ligand showing chromophoric nature with the given metal ions and acts as chemosensor.

Table 1. FT-IR analysis of chemosensor and samples

No.	Sample / Compound	Frequencies, cm ⁻¹	Assigned group
1.	Chemosensor	3623, 3750	N-H
		2982	C-H (aromatic)
		1700	C=O
		1647, 1546	C=C (aromatic)
		1280	C-C
2.	Chemosensor after reacting with 1% Cu.	1046	C-N
		2946	C-H
		2876	C-H (Aliphatic)
		1746	C=O
		1656,	C=C (Aromatic)
3.	Chemosensor after reacting with 1% Hg.	1621,1546	N-H
		3231, 3456	
		2972	C-H (Aliphatic)
		1748	C=O
		1648	C=C (Aromatic)
4.	Chemosensor after reacting with 1% Ni	1195	C-N
		1317	C-O
		3234-3496	N-H
		951	N-Ni ²⁺
		2892	C-H (Aromatic)
		3365	N-H




Figure 1. Amide based chemosensor forms coordination complex with metal ions




Figure 2. UV analysis of Co^{2+} dilutions with chemosensor (100-1000 ppm)

Conclusion

We developed a simple, efficient and eco-friendly method for the synthesis of amide-based chemosensor from commercially available material. We studied the application of chemosensor for qualitative and quantitative analysis of metal ions in a various extracted solution using UV-VIS spectroscopy techniques. This chemosensor helps in the preliminary analysis and detection of Hg^{2+} , Co^{2+} , Ni^{2+} , Cu^{2+} metal ions in aqueous and spiked solutions. This chemosensor has excellent application in forensic chemistry and toxicology to detect the metal poisoning cases.

Acknowledgment

We are thankful to the Department of Forensic Chemistry and Toxicology, Government Institute of Forensic Science, Aurangabad for the permission to use laboratory and sophisticated instrumentation facility. Authors are especially thankful to Dr. S. G. Gupta (Director, Government Institute of Forensic Science, Aurangabad) for support and permitting us to use the sophisticated instrumentation facility. □

References

- Lee, S., Choi, I., Hong, S., Yang, Y.I., Lee, J., Kang, T., Yi, J., Highly selective detection of Cu^{2+} utilizing specific binding between Cu-demethylated superoxide dismutase 1 and the Cu^{2+} ion via surface plasmon resonance spectroscopy, *Chem. Commun.*, **2009**, 41, 6171-6173. doi: 10.1039/b910666a
- Zhang, X., Xiao, Y., Qian, X., A Ratiometric Fluorescent Probe Based on FRET for Imaging Hg^{2+} Ions in Living Cells, *Angew. Chem. Int. Ed.*, **2008**, 47, 8025. doi.org/10.1002/anie.200803246
- Li, L. Q., A Colorimetric and Fluorescent Chemosensor for Cu^{2+} Detection in Aqueous Solution, *Synth. React Inorg. Metal-Org. and Nano-Metal Chem.*, **2016**, 46, 1854-1856. doi.org/10.1080/15533174.2015.1137077
- Que, E. L., Domaille, D. W., Chang, C. J., *Metals in neurobiology: Probing their chemistry and biology with molecular imaging*, *Chem Rev*, **2008**, 108, 1517-1548. doi: 10.1021/cr078203u
- Feng, J., Shao, X., Shang, Z., Chao, J., Wang, Y., Jin, W., A new biphenylcarbonitrile based fluorescent sensor for Zn^{2+} ions and application in living cells, *Chem. Res. Chin. Univ.*, **2017**, 33, 695-701. doi: 10.1007/s40242-017-7084-z
- Patra, C., Bhanja, A. K., Sen, C., Ojha, D., Chattopadhyay, D., Mahapatra, A., Sinha, C., Imine-functionalized thioether Zn(II) turn-on fluorescent sensor and its selective sequential logic operations with H_2PO_4^- , DFT computation and live cell imaging, *RSC Adv.*, **2016**, 6, 53378-53388. doi: 10.1039/C6RA07089B
- Bertini, I., Gray, H. B., Lippard, S. J., Valentin, J. S., *Bioinorg. Chem.*, **1994**, University Science Books, Mill Valley, CA (USA). ISBN 0-935702-57-1
- Park, G. J., Na, Y. J., Jo, H. Y., Lee, S. A., Kim, C., A colorimetric organic chemo-sensor for Co^{2+} in a fully aqueous environment, *Dalton Trans.*, **2014**, 43, 6618-6622. doi: 10.1039/C4DT00423J
- Sun, J., Ye, B., Xia, G., Wang, H. M., A multi-responsive squaraine-based "turn on" fluorescent chemosensor for highly sensitive detection of Al^{3+} , Zn^{2+} and Cd^{2+} in aqueous media and its biological application, *Sens. Actuators B: Chem.*, **2016**, 223, 234-240. doi.org/10.1016/j.snb.2017.03.134
- Liu, X., Lin, Q., Wie, T.-B., Zhang, Y.-M., A highly selective colorimetric chemosensor for detection of nickel ions in aqueous solution, *New J. Chem.*, **2014**, 38, 1418-1423. doi: 10.1039/C3NJ01403G
- Shahabuddin, M., Ashfaq, A. B., Asif, A. B., A new calix[4]arene Schiff base sensor for Hg^{2+} and Au^{3+} , *J. Iran. Chem. Soc.*, **2016**, 13, 2275-2282. https://doi.org/10.1007/s13738-016-0946-3
- Yao, G., Guowen, M., Meiling, W., Qing, H., Chuhong, Z., Zhulin, H., R6G/8-AQ co-functionalized $\text{Fe}_3\text{O}_4@/\text{SiO}_2$ nanoparticles for fluorescence detection of trace Hg^{2+} and Zn^{2+} in aqueous solution, *Sci. China Mater*, **2015**, 58, 550-558. doi: 10.1007/s40843-015-0071-0
- Ji, Y. K., Yun, J. J., Yoon, J. L., Kwan, M. K., Mi, S. S., Wonwoo, N., Juyoung, Y., A Highly Selective Fluorescent Chemosensor for Pb^{2+} , *J. Am. Chem. Soc.*, **2005**, 127, 10107-10111. doi: 10.1021/ja051075b
- Zhou, Y., Zhu, C.-Y., Gao, X.-S., You, X.-Y., Yao, C., Hg^{2+} -Selective Ratiometric and "Off-On" Chemosensor Based on the Azadiene-Pyrene Derivative, *Org. Lett.*, **2010**, 12, 2566-2569. doi: 10.1021/ol1007636
- Liu, W., Xu, L., Sheng, R., Wang, P., Li, H., Wu, S., A Water-Soluble "Switching On" Fluorescent Chemosensor of Selectivity to Cd^{2+} , *Org. Lett.*, **2007**, 9, 3829-3832. doi: 10.1021/ol701620h
- Nadeem, S., Shah, M. R., Khan, B., Hoda, N., Topel, Ö., Supramolecular chemosensor for selective detection of iron in aqueous medium, *Supramol. Chem.*, **2013**, 25, 798-805. doi.org/10.1080/10610278.2013.804186
- Eftekhari-Sis, B., Rezazadeh, Z., Akbari, A., Amini, M., 8-Hydroxyquinoline Functionalized Graphene Oxide: an Efficient Fluorescent Nanosensor for Zn^{2+} in Aqueous Media, *J. Fluoresc.*, **2018**, 28, 1173-1180. doi.org/10.1007/s10895-018-2281-9.
- Yang, Q.-Z., Wu, L.-Z., Zhang, H., Chen, B., Wu, Z. X., Zhang, L. P., Tung, C.-H., A Luminescent Chemosensor with Specific Response for Mg^{2+} , *Inorg. Chem.*, **2004**, 43, 5195-5197. doi: 10.1021/ic049815p
- Wang, M., Liu, X., Lu, H., Wang, H., Qin, Z., Highly Selective and Reversible Chemosensor for Pd^{2+} Detected by Fluorescence, Colorimetry, and Test Paper, *ACS Appl. Mater. Interfaces*, **2015**, 7, 1284-1289. doi: 10.1021/am507479m
- Coskun, A., Akkaya, E. U., Ion Sensing Coupled to Resonance Energy Transfer: A Highly Selective and Sensitive Ratiometric Fluorescent Chemosensor for $\text{Ag}(\text{I})$ by a Modular Approach, *J. Am. Chem. Soc.*, **2005**, 127, 10464-10465. doi: 10.1021/ja052574f
- Lin, Q., Fu, Y.-P., Pei, C., Wei, T. B., Zhang, Y.-M., Rational design, green synthesis of reaction-based dual-channel chemosensors for cyanide anion, *Tetrahedron Letters*, **2013**, 54, 5031-5034. doi.org/10.1016/j.tetlet.2013.07.022

- ²²Parsaee, Z., Haratipour, P., Lariche, M. J., Vojood, A., A novel high-performance nano chemosensor for copper (II) ion based on an ultrasound-assisted synthesized diphenylamine-based Schiff base: Design, fabrication and density functional theory calculations, *Ultrasonics Sonochem.*, **2018**, *41*, 337-349. doi.org/10.1016/j.ultsonch.2017.09.054
- ²³Li, H., Shao, F. Q., Huang, H., Feng, J.-J., Wang, A.-J., Eco-friendly and rapid microwave synthesis of green fluorescent graphitic carbon nitride quantum dots for vitro bioimaging, *Sensors Actuators B*, **2016**, *226*, 506-511. doi.org/10.1016/j.snb.2015.12.018
- ²⁴Basu, A., Suryawanshi, A., Kumawat, B., Dandia, A., Guin, D., Ogale, S. B., Starch (Tapioca) to carbon dots: an efficient green approach to an on-off-on photoluminescence probe for fluoride ion sensing, *Analyst*, **2015**, *140*, 1837-1841. [doi:10.1039/C4AN02340D](https://doi.org/10.1039/C4AN02340D)
- ²⁵Lieberzeit, P. A., Dickert, F. L., Chemosensors in environmental monitoring: challenges in ruggedness and selectivity, *Anal. Bioanal. Chem.*, **2009**, *393*, 467. DOI:10.1007/s00216-008-2464-3
- ²⁶Yilmaz, E., Garipcan, B., Patra, H. K., Uzun, L., Molecular Imprinting Applications in Forensic Science, *Sensors*, **2017**, *17*, 691-695. <https://doi.org/10.3390/s17040691>
- ²⁷Jung, H.-S., Ko, K. C., Kim, G.-H., Lee, A.-R., Na, Y.-C., Kang, C., Lee, J. Y., Kim, J. S., Coumarin-Based Thiol Chemosensor: Synthesis, Turn-On Mechanism, and Its Biological Application, *Org. Lett.*, **2011**, *13*, 1498-1501. <https://doi.org/10.1021/ol2001864>
- ²⁸Bang, J. H., Suslick, K. S., Applications of Ultrasound to the Synthesis of Nanostructured Materials, *Adv. Mater.*, **2010**, *22*, 1039-1059. doi.org/10.1002/adma.200904093

This paper was presented at the “*International Symposium on Exploring New Horizons in Chemical Sciences*”, January 10–12, **2019**, Aurangabad, India (ENHCS–2019).

Received: 11.03.2019

Accepted: 14.07.2019.