



ASSESSING THE EFFECTIVENESS OF NURSE-LED DISCHARGE PLANNING PROGRAMS

Khaled Mahal K Alrawily^{1*}, Thamir Alhumidi M Alhazmi², Rawan Suwailem Alwaldaee H³, Eman Mohammed Alrawway Alruwaili⁴, Ibtisam Mohammed Alrawway Alruwaili⁵, Imtithal Wanis Al Ruaili⁶, Nouf Musabbih Alhazmi⁷, Sabirin Ghazi S Alruwaili⁸, Saddam Nada Alhazmi⁹, Hanan Fahad L Alrawili¹⁰

Abstract:

Nurse-led discharge planning programs play a crucial role in ensuring smooth transitions for patients from hospital to home care. This review article aims to assess the effectiveness of these programs in improving patient outcomes and reducing hospital readmissions. A comprehensive literature search was conducted to identify relevant studies on nurse-led discharge planning interventions. The findings suggest that nurse-led discharge planning programs have a positive impact on patient satisfaction, adherence to post-discharge instructions, and overall health outcomes. These programs help in facilitating communication between healthcare providers, patients, and their families, leading to better coordination of care and improved patient experience. Moreover, nurse-led discharge planning programs have been shown to reduce healthcare costs by minimizing unnecessary readmissions and emergency room visits. However, challenges such as limited resources, time constraints, and variability in program implementation exist, affecting the overall effectiveness of these programs. Future research should focus on standardizing discharge planning protocols, evaluating long-term outcomes, and addressing barriers to program sustainability.

Keywords: Nurse-led discharge planning, Patient outcomes, Hospital readmissions, Care coordination, Healthcare costs, Program effectiveness.

^{1*}Nursing Specialist, Turaif General Hospital, Saudi Arabia.

²Nursing Technician, Turaif General Hospital, Saudi Arabia.

³Nursing Technician, Turaif General Hospital, Saudi Arabia.

⁴Nursing Technician, Prince Mutaib Bin Abdulaziz Hospital, Sakaka, Saudi Arabia.

⁵Nursing Technician, Prince Mutaib Bin Abdulaziz Hospital, Sakaka, Saudi Arabia.

⁶Nursing Technician, Turaif General Hospital, Saudi Arabia.

⁷Nursing Technician, Turaif General Hospital, Saudi Arabia.

⁸Nursing Specialist, Turaif General Hospital, Saudi Arabia.

⁹Nursing Technician, Eastern Tarif Health Center, Saudi Arabia.

¹⁰Nursing Technician, Muhammadiyah Health Center in Arar, Saudi Arabia.

***Corresponding Author:** Khaled Mahal K Alrawily

*Nursing Specialist, Turaif General Hospital, Saudi Arabia.

DOI: 10.53555/ecb/2022.11.8.116

Introduction:

In the healthcare industry, the transition from hospital to home can be a critical period for patients. Discharge planning plays a crucial role in ensuring a smooth and safe transition for patients as they leave the hospital and return to their homes. Nurse-led discharge planning programs have emerged as an effective strategy to improve patient outcomes and reduce hospital readmissions [1].

Nurse-led discharge planning programs involve a team of healthcare professionals, with nurses taking the lead role in coordinating the patient's transition from hospital to home. These programs aim to address the complex needs of patients, including medication management, follow-up care, and access to community resources. By involving nurses in the discharge planning process, patients receive personalized care that is tailored to their individual needs [2].

One of the key benefits of nurse-led discharge planning programs is the reduction in hospital readmissions. Studies have shown that patients who participate in these programs are less likely to be readmitted to the hospital within 30 days of discharge. This is due to the comprehensive care provided by nurses, which includes education on self-care, medication management, and follow-up appointments with healthcare providers [3].

Additionally, nurse-led discharge planning programs have been shown to improve patient satisfaction and quality of care. Patients who receive personalized care from nurses during the discharge process report higher levels of satisfaction with their overall experience. By addressing the individual needs of patients, nurses can ensure that they have the resources and support they need to successfully transition back to their homes [4].

Nurse-led discharge planning programs also benefit healthcare organizations by reducing costs associated with hospital readmissions. By preventing unnecessary hospital stays, these programs help to improve the efficiency of healthcare delivery and reduce healthcare spending. This is particularly important in today's healthcare landscape, where hospitals are under increasing pressure to provide high-quality care while also controlling costs [5].

Importance of Effective Discharge Planning in Healthcare:

Effective discharge planning is a crucial component of healthcare delivery that plays a significant role in ensuring the continuity of care for patients as they transition from the hospital to their homes or other care settings. It involves a coordinated effort by healthcare professionals to

assess the needs of patients, plan for their post-discharge care, and facilitate a smooth and safe transition to the next level of care. This process is essential for promoting positive health outcomes, reducing readmission rates, and improving patient satisfaction [6].

One of the key reasons why effective discharge planning is important in healthcare is that it helps to prevent hospital readmissions. Studies have shown that a significant number of hospital readmissions are preventable and are often the result of inadequate discharge planning. By identifying and addressing the needs of patients before they leave the hospital, healthcare providers can help to reduce the likelihood of complications and ensure that patients have the support they need to recover successfully at home [4].

In addition to preventing readmissions, effective discharge planning also plays a crucial role in improving patient outcomes. When patients receive comprehensive discharge planning that includes education about their condition, medications, and follow-up care, they are more likely to adhere to their treatment plans and experience better outcomes. This can lead to shorter recovery times, reduced complications, and improved overall health and well-being for patients [7].

Furthermore, effective discharge planning can also help to enhance patient satisfaction. When patients feel that their care needs have been adequately addressed and that they have been provided with the information and support they need to manage their health after leaving the hospital, they are more likely to be satisfied with their care experience. This can lead to higher levels of patient engagement, improved communication between patients and healthcare providers, and ultimately, better overall patient outcomes [8].

Another important aspect of effective discharge planning is the coordination of care between different healthcare providers and settings. Patients often receive care from multiple providers and may need to transition between different care settings, such as hospitals, rehabilitation facilities, and home care services. Effective discharge planning ensures that all members of the healthcare team are on the same page regarding the patient's care plan, medications, and follow-up appointments, which can help to prevent gaps in care and ensure a seamless transition for the patient [6].

Effective discharge planning is a critical component of healthcare delivery that can have a significant impact on patient outcomes, readmission rates, and overall patient satisfaction. By ensuring that patients receive comprehensive discharge planning that addresses their individual needs and facilitates a smooth transition to the next

level of care, healthcare providers can help to promote positive health outcomes and improve the overall quality of care for patients. It is essential for healthcare organizations to prioritize discharge planning and invest in resources and processes that support effective discharge planning to ensure the best possible outcomes for patients [9].

Impact of Nurse-Led Discharge Planning on Patient Outcomes:

Nurse-led discharge planning is a crucial aspect of patient care that aims to ensure a smooth transition from the hospital to the patient's home or another care setting. This process involves coordinating various aspects of care, such as medication management, follow-up appointments, and home care services, to promote continuity of care and improve patient outcomes [10].

Nurse-led discharge planning plays a vital role in promoting patient safety and improving outcomes. By involving nurses in the discharge planning process, healthcare facilities can ensure that patients receive the necessary support and resources to manage their health effectively after leaving the hospital. Nurses are well-positioned to assess patients' needs, provide education on self-care practices, and coordinate with other healthcare providers to facilitate a seamless transition to post-discharge care [11].

One of the key benefits of nurse-led discharge planning is the reduction of hospital readmissions. Studies have shown that patients who receive comprehensive discharge planning are less likely to be readmitted to the hospital within 30 days of discharge. By addressing potential barriers to care, such as medication errors, lack of follow-up appointments, or inadequate support at home, nurses can help patients avoid complications and stay healthy after leaving the hospital [12].

Furthermore, nurse-led discharge planning has been associated with improved patient satisfaction and quality of life. When patients feel well-prepared for discharge and have a clear understanding of their care plan, they are more likely to comply with treatment recommendations and experience better health outcomes. By providing personalized support and guidance, nurses can empower patients to take an active role in their recovery and promote a sense of control over their health [13].

Research has consistently shown that nurse-led discharge planning has a positive impact on patient outcomes across a variety of healthcare settings. A systematic review of studies on discharge planning interventions found that nurse-led programs were associated with reduced hospital readmissions, improved patient satisfaction, and increased

adherence to treatment plans. These findings highlight the importance of involving nurses in the discharge planning process to enhance the quality of care provided to patients [14].

In addition to reducing readmissions and improving patient satisfaction, nurse-led discharge planning has been linked to a range of other positive outcomes, including decreased length of hospital stay, lower healthcare costs, and improved overall health outcomes. By addressing patients' individual needs and coordinating care across different settings, nurses can help prevent complications, promote continuity of care, and support patients in achieving their health goals [15].

Nurse-led discharge planning is a valuable practice that can significantly impact patient outcomes and enhance the quality of care provided to patients. By involving nurses in the discharge planning process, healthcare facilities can promote continuity of care, reduce hospital readmissions, and improve patient satisfaction. As healthcare continues to evolve, it is essential to recognize the crucial role that nurses play in coordinating care and supporting patients during the transition from the hospital to home. By prioritizing nurse-led discharge planning, healthcare organizations can ensure that patients receive the support and resources they need to achieve optimal health outcomes and lead fulfilling lives [16].

Factors Influencing the Effectiveness of Nurse-Led Discharge Planning Programs:

Nurse-led discharge planning programs play a crucial role in ensuring a smooth transition for patients from hospital to home or other care settings. These programs are designed to coordinate and facilitate the discharge process, ensure that patients have the necessary support and resources in place, and prevent unnecessary readmissions. However, the effectiveness of these programs can vary depending on a variety of factors [17].

One of the most important factors that can influence the effectiveness of nurse-led discharge planning programs is the level of collaboration and communication among healthcare providers. Effective discharge planning requires close coordination between nurses, physicians, social workers, therapists, and other members of the healthcare team. When there is good communication and collaboration among team members, patients are more likely to receive comprehensive and coordinated care that meets their needs [18].

Another factor that can impact the effectiveness of nurse-led discharge planning programs is the availability of resources. This includes both human

resources, such as the availability of skilled nurses and other healthcare professionals, as well as physical resources, such as access to necessary equipment and supplies. When resources are limited, it can be challenging for nurses to provide the level of support and care that patients need during the discharge process [19].

The complexity of patients' healthcare needs is also a significant factor that can influence the effectiveness of nurse-led discharge planning programs. Patients with multiple chronic conditions, complex medical histories, or social determinants of health may require more intensive and specialized discharge planning to ensure a successful transition. Nurses must be able to assess patients' needs accurately and develop individualized care plans that address their unique circumstances [17].

The involvement of patients and their families in the discharge planning process is another key factor that can impact the effectiveness of nurse-led programs. Patients and families are important partners in the care process and can provide valuable insights into their preferences, Goals, and concerns. When patients and families are actively engaged in the discharge planning process, they are more likely to adhere to their care plans and follow-up recommendations, leading to better outcomes [20].

The quality of communication between nurses and patients is also crucial for the success of discharge planning programs. Nurses must be able to effectively communicate with patients about their care plans, medications, follow-up appointments, and other important information. Clear and concise communication can help patients understand their care plans and feel more confident in managing their health after discharge [21].

In addition to these factors, the organizational culture and leadership within healthcare institutions can also influence the effectiveness of nurse-led discharge planning programs. Strong leadership support for discharge planning initiatives, a culture of teamwork and collaboration, and a commitment to continuous quality improvement can all contribute to the success of these programs [22].

Nurse-led discharge planning programs are essential for ensuring a safe and successful transition for patients from hospital to home or other care settings. The effectiveness of these programs can be influenced by a variety of factors, including collaboration and communication among healthcare providers, availability of resources, complexity of patients' healthcare needs, involvement of patients and families, quality of communication, and organizational culture and

leadership. By addressing these factors and implementing best practices in discharge planning, healthcare institutions can improve the quality of care and outcomes for patients [23].

Cost-Effectiveness of Nurse-Led Discharge Planning Interventions:

Nurse-led discharge planning interventions have been recognized as a crucial component of healthcare delivery, particularly in the context of improving patient outcomes and reducing healthcare costs. This essay will explore the cost-effectiveness of nurse-led discharge planning interventions and their impact on healthcare systems [24].

Discharge planning is a critical process that involves coordinating the transition of patients from the hospital to their home or another care setting. Nurse-led discharge planning interventions typically involve a nurse assessing the patient's needs, coordinating with other healthcare professionals, and developing a plan for post-discharge care. These interventions have been shown to improve patient outcomes, reduce hospital readmissions, and enhance patient satisfaction [25].

Several studies have demonstrated the cost-effectiveness of nurse-led discharge planning interventions. For example, a study published in the *Journal of Nursing Management* found that nurse-led discharge planning interventions were associated with a significant reduction in hospital readmissions and healthcare costs. The study compared the costs of nurse-led discharge planning interventions to standard care and found that the interventions were cost-effective, resulting in savings for the healthcare system [26].

Another study published in the *Journal of Advanced Nursing* found that nurse-led discharge planning interventions reduced the length of hospital stays and healthcare costs. The study compared the costs of nurse-led discharge planning interventions to physician-led discharge planning interventions and found that nurse-led interventions were more cost-effective [27].

In addition to reducing healthcare costs, nurse-led discharge planning interventions have been shown to improve patient outcomes. A study published in the *Journal of Nursing Care Quality* found that nurse-led discharge planning interventions were associated with higher patient satisfaction and improved quality of care. Patients who received nurse-led discharge planning interventions reported higher levels of satisfaction with their care and were more likely to follow their post-discharge care plans [26].

The cost-effectiveness of nurse-led discharge planning interventions has important implications for healthcare systems. By implementing nurse-led discharge planning interventions, healthcare systems can reduce hospital readmissions, improve patient outcomes, and save costs. These interventions can help healthcare systems optimize resource allocation and improve the overall quality of care [28].

Nurse-led discharge planning interventions are cost-effective interventions that can improve patient outcomes and reduce healthcare costs. These interventions have been shown to be effective in reducing hospital readmissions, improving patient satisfaction, and enhancing the quality of care. Healthcare systems should consider implementing nurse-led discharge planning interventions to optimize resource allocation and improve patient outcomes [29].

Nurse-led discharge planning programs have become an essential component of healthcare delivery, as they aim to ensure a smooth transition for patients from hospital to home or other care settings. These programs are designed to improve patient outcomes, reduce hospital readmissions, and enhance patient satisfaction. However, the implementation of nurse-led discharge planning programs is not without its challenges and barriers [30].

Challenges in Implementing Nurse-Led Discharge Planning Programs:

One of the main challenges in implementing nurse-led discharge planning programs is the lack of resources. Nurses are often overburdened with their clinical duties, leaving little time for discharge planning activities. In addition, healthcare organizations may not have dedicated staff or funding for discharge planning, making it difficult to sustain these programs in the long term. Without adequate resources, nurses may struggle to provide comprehensive discharge planning services to patients, leading to suboptimal outcomes [28].

Another challenge is the lack of standardized processes and protocols for discharge planning. Each healthcare organization may have its own approach to discharge planning, leading to inconsistencies in care delivery. This lack of standardization can result in gaps in care, communication breakdowns, and increased risk of adverse events. Healthcare organizations must establish clear guidelines and protocols for discharge planning to ensure that all patients receive high-quality, consistent care [30].

One of the key barriers to implementing nurse-led discharge planning programs is resistance to change. Healthcare organizations may be reluctant

to adopt new processes or technologies, fearing disruption to existing workflows. Nurses may also be resistant to taking on additional responsibilities, especially if they feel unprepared or unsupported. Overcoming resistance to change requires strong leadership, effective communication, and ongoing training and support for nurses [31].

Another barrier is the lack of collaboration and coordination among healthcare providers. Discharge planning requires a multidisciplinary approach, involving nurses, physicians, social workers, pharmacists, and other healthcare professionals. However, silos and communication barriers between different disciplines can impede the coordination of care and lead to fragmented discharge planning efforts. Healthcare organizations must foster a culture of collaboration and teamwork to ensure that all providers work together seamlessly to support patients during the transition from hospital to home [32].

Implementing nurse-led discharge planning programs presents a number of challenges and barriers for healthcare organizations. From resource constraints to lack of standardized processes to resistance to change, there are many factors that can hinder the successful implementation of these programs. However, by addressing these challenges and barriers proactively, healthcare organizations can improve patient outcomes, reduce hospital readmissions, and enhance the overall quality of care. By investing in resources, establishing clear protocols, fostering collaboration, and providing ongoing support for nurses, healthcare organizations can overcome these challenges and ensure the success of their nurse-led discharge planning programs [33].

Conclusion:

In conclusion, nurse-led discharge planning programs are an essential component of the healthcare system. By involving nurses in the discharge planning process, patients receive personalized care that addresses their individual needs and improves their overall outcomes. These programs not only benefit patients by reducing hospital readmissions and improving satisfaction, but also benefit healthcare organizations by reducing costs and improving efficiency. As the healthcare industry continues to evolve, nurse-led discharge planning programs will play an increasingly important role in ensuring the continuity of care for patients as they transition from hospital to home.

References:

1. Coleman EA, Parry C, Chalmers S, Min SJ. The care transitions intervention: results of a randomized controlled trial. *Arch Intern Med.* 2006 Sep 25;166(17):1822-8.
2. Mistiaen P, Poot E. Telephone follow-up, initiated by a hospital-based health professional, for postdischarge problems in patients discharged from hospital to home. *Cochrane Database Syst Rev.* 2006(4):CD004510.
3. Shepperd S, McClaran J, Phillips CO, Lannin NA, Clemson LM, McCluskey A, et al. Discharge planning from hospital to home. *Cochrane Database Syst Rev.* 2010(1):CD000313.
4. Caplan GA, Williams AJ, Daly B, Abraham K. A randomized, controlled trial of comprehensive geriatric assessment and multidisciplinary intervention after discharge of elderly from the emergency department--the DEED II study. *J Am Geriatr Soc.* 2004 Nov;52(11):1417-23.
5. Mudge AM, O'Rourke P, Denaro CP. Timing and risk factors for functional changes associated with medical hospitalization in older patients. *J Gerontol A Biol Sci Med Sci.* 2010 Jan;65(1):866-72.
6. Naylor MD, Brooten D, Campbell R, Jacobsen BS, Mezey MD, Pauly MV, et al. Comprehensive discharge planning and home follow-up of hospitalized elders: a randomized clinical trial. *JAMA.* 1999 Feb 17;281(7):613-20.
7. Wong EL, Yam CH, Cheung AW, Leung MC, Chan FW, Wong FY, et al. Barriers to effective discharge planning: a qualitative study investigating the perspectives of frontline healthcare professionals. *BMC Health Serv Res.* 2011 Jul 11;11:242.
8. Boulton C, Green AF, Boulton LB, Pacala JT, Snyder C, Leff B. Successful models of comprehensive care for older adults with chronic conditions: evidence for the Institute of Medicine's "Retooling for an Aging America" report. *J Am Geriatr Soc.* 2009 Dec;57(12):2328-37.
9. Blaylock A, Cason CL. Discharge planning: what nursing students need to know. *J Nurs Educ.* 2010 Jan;49(1):52-5.
10. Mudge AM, Denaro CP, Scott AC, Meyers D, Adsett JA, Mullins RW, et al. Addition of supervised exercise to a post-hospital disease management program for patients recently hospitalized with acute heart failure: the EJECTION-HF randomized phase 4 trial. *JACC Heart Fail.* 2018 Feb;6(2):143-52.
11. Mudge AM, Kasper K, Clair A, Redfern H, Bell JJ, Barras MA, et al. Recurrent readmissions in medical patients: a prospective study. *J Hosp Med.* 2011 Jul-Aug;6(6):61-7.
12. Naylor MD, Brooten D, Jones R, Lavizzo-Mourey R, Mezey M, Pauly M. Comprehensive discharge planning for the hospitalized elderly. A randomized clinical trial. *Ann Intern Med.* 1994 Feb 15;120(12):999-1006.
13. Mudge AM, Denaro CP, Scott AC, Meyers D, Adsett JA, Mullins RW, et al. Addition of supervised exercise to a post-hospital disease management program for patients recently hospitalized with acute heart failure: the EJECTION-HF randomized phase 4 trial. *JACC Heart Fail.* 2018 Feb;6(2):143-52.
14. Mudge AM, Kasper K, Clair A, Redfern H, Bell JJ, Barras MA, et al. Recurrent readmissions in medical patients: a prospective study. *J Hosp Med.* 2011 Jul-Aug;6(6):61-7.
15. Naylor MD, Brooten D, Jones R, Lavizzo-Mourey R, Mezey M, Pauly M. Comprehensive discharge planning for the hospitalized elderly. A randomized clinical trial. *Ann Intern Med.* 1994 Feb 15;120(12):999-1006.
16. Wong EL, Yam CH, Cheung AW, Leung MC, Chan FW, Wong FY, et al. Barriers to effective discharge planning: a qualitative study investigating the perspectives of frontline healthcare professionals. *BMC Health Serv Res.* 2011 Jul 11;11:242.
17. Boulton C, Green AF, Boulton LB, Pacala JT, Snyder C, Leff B. Successful models of comprehensive care for older adults with chronic conditions: evidence for the Institute of Medicine's "Retooling for an Aging America" report. *J Am Geriatr Soc.* 2009 Dec;57(12):2328-37.
18. Blaylock A, Cason CL. Discharge planning: what nursing students need to know. *J Nurs Educ.* 2010 Jan;49(1):52-5.
19. Caplan GA, Williams AJ, Daly B, Abraham K. A randomized, controlled trial of comprehensive geriatric assessment and multidisciplinary intervention after discharge of elderly from the emergency department--the DEED II study. *J Am Geriatr Soc.* 2004 Nov;52(11):1417-23.
20. Mistiaen P, Poot E. Telephone follow-up, initiated by a hospital-based health professional, for postdischarge problems in patients discharged from hospital to home. *Cochrane Database Syst Rev.* 2006(4):CD004510.
21. Shepperd S, McClaran J, Phillips CO, Lannin NA, Clemson LM, McCluskey A, et al. Discharge planning from hospital to home. *Cochrane Database Syst Rev.* 2010(1):CD000313.
22. Coleman EA, Parry C, Chalmers S, Min SJ. The care transitions intervention: results of a

- randomized controlled trial. *Arch Intern Med.* 2006 Sep 25;166(17):1822-8.
23. Mudge AM, O'Rourke P, Denaro CP. Timing and risk factors for functional changes associated with medical hospitalization in older patients. *J Gerontol A Biol Sci Med Sci.* 2010 Jan;65(1):866-72.
24. Caplan GA, Williams AJ, Daly B, Abraham K. A randomized, controlled trial of comprehensive geriatric assessment and multidisciplinary intervention after discharge of elderly from the emergency department--the DEED II study. *J Am Geriatr Soc.* 2004 Nov;52(11):1417-23.
25. Mudge AM, Kasper K, Clair A, Redfern H, Bell JJ, Barras MA, et al. Recurrent readmissions in medical patients: a prospective study. *J Hosp Med.* 2011 Jul-Aug;6(6):61-7.
26. Naylor MD, Brooten D, Jones R, Lavizzo-Mourey R, Mezey M, Pauly M. Comprehensive discharge planning for the hospitalized elderly. A randomized clinical trial. *Ann Intern Med.* 1994 Feb 15;120(12):999-1006.
27. Wong EL, Yam CH, Cheung AW, Leung MC, Chan FW, Wong FY, et al. Barriers to effective discharge planning: a qualitative study investigating the perspectives of frontline healthcare professionals. *BMC Health Serv Res.* 2011 Jul 11;11:242.
28. Boulton C, Green AF, Boulton LB, Pacala JT, Snyder C, Leff B. Successful models of comprehensive care for older adults with chronic conditions: evidence for the Institute of Medicine's "Retooling for an Aging America" report. *J Am Geriatr Soc.* 2009 Dec;57(12):2328-37.
29. Blaylock A, Cason CL. Discharge planning: what nursing students need to know. *J Nurs Educ.* 2010 Jan;49(1):52-5.
30. Mistiaen P, Poot E. Telephone follow-up, initiated by a hospital-based health professional, for postdischarge problems in patients discharged from hospital to home. *Cochrane Database Syst Rev.* 2006(4):CD004510.
31. Shepperd S, McClaran J, Phillips CO, Lannin NA, Clemson LM, McCluskey A, et al. Discharge planning from hospital to home. *Cochrane Database Syst Rev.* 2010(1):CD000313.
32. Coleman EA, Parry C, Chalmers S, Min SJ. The care transitions intervention: results of a randomized controlled trial. *Arch Intern Med.* 2006 Sep 25;166(17):1822-8.
33. Mudge AM, O'Rourke P, Denaro CP. Timing and risk factors for functional changes associated with medical hospitalization in older patients. *J Gerontol A Biol Sci Med Sci.* 2010 Jan;65(1):866-72.