



**Impact of Agro N.P.K Fertilizer induced Aquatic Pollution and Associated Histopathological Infractions on African Catfish (*Clarias gariepinus*) cultured in Calabar, Nigeria**

**Ettah Ivon. A<sup>1</sup>, Agbor R. B<sup>2</sup>, Osondu Anyanwu C<sup>1</sup>, Inah Simon Alain<sup>3</sup>, Edodi I. O<sup>1</sup> and Godwin M. Ubi<sup>2</sup>**

<sup>1</sup>Department of Science, Laboratory Technology, Faculty of Biological Science, University of Calabar, Calabar, Nigeria.

<sup>2</sup>Department of Genetics and Biotechnology, Faculty of Biological Sciences, University of Calabar, Nigeria

<sup>3</sup>Department of Public Health, Faculty of Allied Medical Sciences, University of Calabar, Calabar, Nigeria

ORCID ID: 0000-0003-0424-234X

SCOPUS Author ID:57214818007

[ubiology.gu@gmail.com](mailto:ubiology.gu@gmail.com)

**ABSTRACT**

Toxicity of Agro NPK fertilizer on the histopathology of the African catfish (*C. gariepinus*) sub adult was investigated in Calabar. The experiment was conducted in triplicates of four treatments. The fish were exposed to concentrations 1.0 g/L, 2.0 g/L, 3.0 g/L and 4.0 g/L. Behavioral changes in fish exposed to different concentrations of Agro NPK fertilizer ranged from erratic swimming, moribund movement, jumping and lack of balance. Similar changes were not observed in the control throughout the experiment. At 1.0g/l concentrations, the mean mortality was 80% with an initial erratic swimming. It was observed that mean mortality increased progressively with increase in concentration of the pollutant (Agro NPK fertilizer). At concentration 3.0 g/L, mean mortality was observed within 72 hours of exposure preceding moribund swimming while the highest mortality was observed at 4.0 g/L. The concentration at which 50% of the experimental fish were killed (LC<sub>50</sub>) was 1.9 g/L. Results from statistical analysis indicated that mortality varied ( $p < 0.05$ ) significantly with concentrations as higher concentrations recorded higher mortalities. However, mean values showed highest mortality (74%) with Agro NPK fertilizer. Histopathology of the liver of fish exposed to the Agro NPK fertilizer after 96 hours revealed severe damage and necrosis of the liver cells when compared with the control which had a normal distribution of hepatic cells. Similarly, results from histopathology of the gills of the exposed fish to Agro NPK fertilizer revealed fusion and damages to gills lamellae when compared with the control which has normal distribution of the gills lamellae. Severe changes in the epithelia of the gill arch were also observed. Findings from histopathology of the intestine also revealed severe damages to the intestinal walls and mucosa when compared with the control. And in the same vein, histopathology result on the effect of Agro NPK fertilizer on the intestine of the exposed fish revealed chronic damages to the intestinal mucosa when viewed with AMSCOPE electron microscope at x10 magnification. It was therefore concluded that Agro NPK fertilizer causes significant and severe damages to *C. gariepinus* and recommended the use of ring method other than the broadcasting method of application of Agro NPK fertilizer on farmlands which often results to run off to aquatic bodies to reduce the negative impact on the histopathology of aquatic organisms.

**Keywords:** Agro NPK fertilizer, aquatic pollution, *Clarias gariepinus*, Liver and Gills damage, Mortality

## 1. Introduction

African catfish (*Clarias gariepinus*) are sharp tooth catfish, eel-like in nature, usually dark gray or black with coloration on the back, fading to a white belly. In Africa, the catfish has been reported as being second in size only to the Vundu of the Zambesian waters (Roosendaal, *et al.*, 2011). It has an average adult length of 1.0-1.5m. It reaches maximum length of 1.7m and can weigh up to 60kg. These fish have slender bodies, flat bony heads, notably flatter in the genus *Silurus*, and broad terminate mouths with four pairs of barbels. They also have large accessory breathing organs composed of modified gill arches and only the pectoral fins have spines. African catfish is a nocturnal fish like many other catfish. It feeds on both living and dead animal matter. Because of its wide mouth, it is able to swallow relatively large prey whole. It is able to crawl on dry ground to escape drying pools and it is also able to survive in shallow mud for long period of time in between raining seasons, (Roosendaal, *et al.*, 2011).

It spawns around inundated areas of river, lakes and streams mostly at nights (Ogundiran, 2011). One of the reasons for the choice of this genus of fish for this research work is because it has been found to be a biomarker in the aquatic environment (Abdel-Monein, 2005). Other reasons for this choice of this fish includes its' hardiness and ability to tolerate adverse water quality conditions, its ability to grow fast and feed on large variety of agriculture by products and its ability to tolerate difficult conditions in captivity (Froese *et al* 2014).

Most chemicals exhibit deleterious effects on the aquatic environment if not properly handled or controlled. Industrial products beneficial to man can also pose serious threats to man and the entire environment (Abbas, *et al.*, 2003). Most of the commonly used industrial products in Calabar, Nigeria include Agro NPK fertilizer, fertilizers, kerosene and a host of others (Ivon *et al.*, 2016). Pollution is a global menace that affects all ecological habitats. It is the introduction of foreign toxic substances capable of causing harm to living organisms in the ecosystem and the entire environment (Adham, *et al.*, 2002).

Pollution occurs either on land, air or water. Man depends heavily on water for domestic, industrial and agricultural uses. Poorer water quality means water pollution (Antonio *etal.*, 2007). No nation is completely free from the global menace of water pollution (Ayoola, 2010). This era of globalization and industrialization has sky rocketed the rates of aquatic pollution due to the increasing volume of industrial and domestic effluents that find their way into fresh water and marine habitats, thus altering the balance of these ecosystems (Adams *et al.*, 1989).

In developing countries like Nigeria, only few chemicals have been ecologically tested for safety in spite of their environmental impacts (Bucher *et al.*, 2013). Every ocean and every continent, from the tropics to the once-pristine polar regions is contaminated (World Wide Fund, 2014). The effects of these activities are often outrageous with corresponding immediate or later consequences to both direct and indirect dependents of these water bodies (Byne, *et al.* 1994). Strict government legislation and heavy penalties have not in any way changed these negative trends of indiscriminate disposal of chemicals into our water bodies. Detergent is one of the most toxic pollutants we have in the world today inflicting so much damages to the environment and the organisms thereon especially those in the aquatic environment (Ndome and Ivon, *et al.* 2016).

Unrestricted municipal NPK fertilizer dissolve in rain water as run off, sewage discharge into water bodies, effluents from chemical industries and direct dumping of biodegradable and non-biodegradable wastes are other sources of water pollution. Apparently, human and ecological disorders experienced in industrial settlements as a result of improper disposal of chemicals such as chemical inorganic fertilizers effluents are alarming (Ewa-Oboho *et al.*, 2007).

Agro NPK fertilizer apart from altering water quality and chemistry, inflicts heavy and sometimes mortal damages to aquatic organisms and their dependents, (Hazari, *et al.*, 2013). However, the fish habitats are being contaminated heavily through a number of aquatic pollutants (Rajathy, 1991). Agro NPK fertilizers are widely used in both subsistence and commercial farmlands and are mostly used in boosting crop growth, reproduction and development (Lenntech, 2014). Most pollutants including Agro NPK fertilizer find their way into water bodies through surface run offs due to broadcasting methods of applications, direct discharge through rains streams, rivers into water bodies (Lenntech, 2015). Agro NPK fertilizer also possess the tendencies of unleashing poisonous effects in all types of aquatic life if found in sufficient quantities and these includes the biodegradable inorganic chemical Agro NPK fertilizer (Lenntech, 2014). Most fishes will die when inorganic chemical fertilizers concentration approach 15 parts per million (Ndome and Ivon, 2016).

Agro NPK fertilizer concentrations as low as 5ppm will kill fish eggs (Mckenzie, *et al.*, 1989). As a result of the enormous adverse effects, Agro NPK fertilizer has on the aquatic environment, they are listed among the major aquatic pollutants in the world (Morah, *et al.*, 2003). Their effect on the environment outweighs any other advantage(s). Its affordability and high efficiency in improving plant growth and reproduction makes a greater number of people prefer applying it to crops in Nigeria (Nalecz-Jawecki, *et al.*, 2003). One of the aquatic organisms often affected drastically by these inorganic chemical fertilizer pollutants is the African catfish (Ojikutu, *et al.*, 2012). Apart from being used for enriching the soil fertility and nutritional status, Agro NPK fertilizer has been found to be very effective for enriching the soil flora.

Agro NPK fertilizers are inorganic compounds, which have both polar and non-polar characteristics. They tend to exist at phase boundaries, where they are associated with both polar and non-polar media.

The Agro NPK fertilizers are used to enhance growth, reproduction and development of cultivated crops (Lenntech 2014). It is in view of the high patronage and usage of this hemical inorganic compound that this research was tailored to unveil the histopathological damages associated with the indiscriminate pollution of aquatic environment with this chemical pollutant.

## **Materials and Methods**

### **2.1. Study Location**

The University of Calabar is located in between the Calabar municipality and Calabar south local government areas of Cross River State, Nigeria. It is bounded to the east by the great quo river, Calabar is the capital of Cross River State, Nigeria. It is located geographically at 40 57” North 80 191 0’ East. Cross River State is one of the States in the Niger Delta, South South Region of Nigeria. The State shares a maritime boundary with Cameroon in the east, AkwaIbom State in the South, Abia and Ebonyi States in the West and Benue State in the North. It is geographically located at 50451N, 80301E /50751N, 8.50E, Holzloener, (2002). The State

according to National Population Commission (NPC) (2005) has a population of 3.2 million people with a land mass of 20, 156 km<sup>2</sup> (7,782sqm).

The temperature in Calabar ranges from 21.050 to 33.150 Celsius. It is often at its maximum within the months of January to April. Rainfall in Calabar is not stable and consistent as it rains through the year. It rains more between the months of April and November with the peak between May and October. Calabar has an average annual precipitation of 2718mm. Holzloener, (2002)

Calabar is situated in the tropical rain forest with mangroves and swamps in some areas. It is described as 'clean and green State'. Most of the original vegetation in the study area has been replaced as a result of agriculture, industrial and building activities (Holzloehner, 2002).

Activities in the rural areas are mainly agriculture and petty trading, the urban areas are characterized by heavy commercial activities, industrialization and tertiary education. Fish farming is also one of the major human activities in both rural and urban areas of the study location (Holzloehner, 2002).

## 2.2 The fish farm

The University of Calabar Fish Farm is located about 1 kilometer away from the research laboratory. The fish farm harbours various sizes and species of fishes ranging from fry, fingerlings sub adults and adults. This research was carried out in the Fish Pathology Laboratory of the Faculty of Oceanography, University of Calabar. Collection of samples and gathering of materials was preceded by a thorough reconnaissance survey.

## 2.3 Sources of the pollutants

The Agro NPK fertilizer was purchased from Ministry of Agriculture and Natural Resources, Calabar in Cross River State, Nigeria.




FIGURE 1: Map of the University of Calabar showing the fish farm

## 2.4 Field investigation

A trip was taken to the university of Calabar fish farm to ascertain the availability of African catfish (*Clarias gariepinus*) sub adults. After confirming the availability, the desired quantity of 600 sub adults were booked.

### 2.4.1 Collection and transportation of the study fish

A total of 600 *C. gariepinus* sub adults with a mean weight of  $8.5 \pm 0.2$  g were purchased from the University of Calabar fish farm. Samples were carefully collected and transferred into a plastic container and transported to the Faculty of Oceanography, University of Calabar, Cross River State, Nigeria. The study fishes were transported to the laboratory in transparent plastic containers by car (Ekanem, 2012).to the laboratory which is about five minutes drive from the fish farm.

### 2.4.2 Laboratory studies

A whole day was used in arranging the laboratory. Unwanted and obstructive materials were removed and the needed aquaria thoroughly washed and dried.

## 2.5 Acclimation of study specimens

The sub adult fish were transferred into a laboratory aquarium (80 x 30 x 30 cm<sup>3</sup>) and allowed to acclimatize in this holding tank in the laboratory condition for one weeks at a temperature  $30.02 \pm 0.09^{\circ}\text{C}$  and a pH of 8, the sub adults were fed once daily with commercial feed (Copen's) at 5% of their body weight. The unconsumed feeds and faeces were removed from the holding tank and the water in the tank was changed every 24 hours as recommended by Reish and Oshida (1987).

## 2.6 Pollutants used for Experiments

The pollutants selected for the experiment is the Agro NPK fertilizer 15:15:15.

## 2.7 Stocking of the test fish

Ten (10) sub adults were carefully introduced into each of the plastic aquaria containing the measured toxicant and another ten (10) introduced to the control aquaria using a hand sieve, Ekanem, (2019). This procedure was repeated for all the experiments.

## 2.8 Range finding tests

Series of range finding tests were conducted using the toxicants and the sub-adult fishes to determine the concentration boundaries before the actual experiment.

## 2.9 Toxicity experiments with Agro NPK fertilizer

The experiment was carried out in triplicates of four treatments i.e. 0.00 g/L (control), 1.0g/L, 2.0g/L and 3.0g/L and 4.0g/L. Ten (10) sub adults of *C. gariepinus* were stocked in each of the four glass aquaria (25 x 15.5 x 15.5 cm<sup>3</sup>) in triplicates for the experiment. The experiment was monitored periodically; observation and responses were taken at intervals of 24, 48, 72 and 96 hour respectively

### 2.10 Histopathology of the gills, liver and intestines of the fish

After the 96 hours exposure to Agro NPK fertilizer toxicant, the gills, liver and intestines of the test and control fishes were removed by dissection and fixed in separate containers containing 10% phosphate buffered formalin (PBF). Samples were allowed to fix for at least 48 hours before they were subjected to manual tissue processing.

Samples were washed thoroughly in water to remove excess fixatives. They were dehydrated through graded series of alcohol, processed into wax, sectioned with a rotary microtome, mounted on glass slides, de-waxed and stained with hematoxylin and eosin stain for microscopic examination at magnification X10, X40 and X100.

### 2.11 Statistical analysis

Data obtained were analyzed using analysis of variances (ANOVA) for significant difference ( $p < 0.05$ ) at 95% confidence limit. Significant treatment means were separated using the least significant difference (LSD) test.

## Results

### 3.1 Necrosis of liver cells of fish exposed to different concentrations of Agro NPK Fertilizer

Plates 1 shows normal distribution of liver cells while plate 2 shows various degrees of necrosis of livers cells at 1.0 and 2.0g/l concentration respectively. Plate 3 shows damages to the hepatic cells of the liver at 3.0g/l concentration of Agro NPK fertilizer when viewed with electron microscope with x 10 magnification. Plate 4 shows serious necrosis of liver cells at 4.0g/l concentration when viewed with an electron microscope at x10 magnification. In all, a comparison of plates 1, 2, 3 and 4 with the control (Plate 5) revealed serious alterations of the liver cells when compared with control.




**Plate 1:** Normal distribution of liver cells




**Plate 2:** Necrotic cells of liver at 1.0 g/l conc.



**Plate 3:** Necrotic cells of the liver at 2.0 g/l conc.




**Plate 4:** Damages to hepatic cells of the liver at 3.0g/l conc. of NPK Fertilizer (x 10)



**Plate 5:** Severe Necrosis of liver cells at 4.0 g/l conc. of NPK Fertilizer

### **3.1.2 Damages to the gill lamellae of fishes exposed to different concentrations of Agro NPK fertilizer**


Plates 6 to 10 show damages to gills lamellae of fish exposed to Agro NPK fertilizer at 1.0g/l to 4.0g/l concentrations respectively when compared to the control (plate 6) which has normal distribution of gill lamellae when viewed with an electron microscope at x 10 magnification. Plates 7 and 8 showed damages to the gill lamellae at 1.0 and 2.0 g/l concentration of Agro NPK fertilizer while plates 9 and 10 shows fusion and severe fusion of gills lamellae of fish exposed to Agro NPK fertilizer at concentration 3.0 and 4.0g/l respectively when view with an electron microscope at x 10 magnification. In all, the effects were severe as the concentration increased.




**Plate 6:** Normal distribution of gill lamellae in control ( x 10)



**Plate 7:** Damage to gill lamellae of fish Expose to 1.0 g/l of NPK Fertilizer



**Plate 8:** Damage to gill lamellae of fish exposed to 3.0 g/l of NPK Fertilizer



**Plate 9:** Fusions of the gill lamellae of fish Exposed to 3.0g/l of NPK fertilizer (x 10).



**Plate 10:** severe fusions of the gill lamellae of fish exposed to 4.0 g/l of NPK fertilizer (x 10)

### **3.1.3 Damages to the intestines and intestinal epithelial cells of fish exposed to different concentrations of NPK fertilizer**

Plates 12 to 15 showed different degrees of damages to the intestinal epithelial cells at concentration of 1.0 to 4.0g/l respectively when viewed with an electron microscope at x10 magnification. Plates 12 and 13 revealed damages to the intestines of the fishes exposed to 1.0 and 2.0 g/l of the Agro NPK fertilizer while plates 14 and 15 revealed damages and serious damages to the intestinal mucosa cells of the test fish respectively. The effects increased with increase in concentration.



**Plate 11** Normal intestine in the control (x 10):




**Plate 12:** Damage to the intestine of fish Exposed to 1.0 g/l of NPK Fertilizer



**Plate 13:** Damage to the intestine of fish exposed to 2.0 g/l of NPK Fertilizer





**Plate 14:** Damage to the intestinal mucosa  
Cells of the intestine of fish exposed to  
3.0 g/l of NPK Fertilizer



**Plate 15:** Severe Damage to the intestinal mucosa  
Cells of the intestine of fish exposed to 4.0 g/l  
of NPK Fertilizer (x 10)

## Discussion

Toxicity of Agro NPK fertilizer on the histopathology of the African catfish (*C. gariepinus*) sub adult was investigated. The experiment was conducted in triplicates of four treatments. Behavioral changes in fish exposed to different concentration of Agro NPK fertilizer ranged from erratic swimming, moribund movement, jumping and lack of balance. Similar changes were not observed in the control throughout the experiment. This observation is similar to that of Ndome *et al.*, (2013).

Agro NPK Fertilizer is composed of linear nitrate, phosphate and potash as active ingredients. The ability of these chemicals to cause behavioral changes in *C. gariepinus* has been reported by Ndome *et al.*, (2010) and Ogundiran, (2010). The fish were exposed to concentrations 1.0g/L, 2.0g/L, 3.0g/L and 4.0g/L. At 1.0g/l concentration, the mean mortality was 80% with an initial erratic swimming. It was observed that mean mortality increased drastically with increase in concentration of the pollutant (NPK fertilizer). At concentration 3.0g/L, total mortality was observed within 72hours of exposure preceding moribund swimming while total mortality was observed in 4.0g/L. The concentration at which 50% of the experimental fish were killed (LC<sub>50</sub>) was 0.9g/L. Results from statistical analysis indicated that mortality varied significantly with concentrations as higher values recorded higher mortalities. However, mean values showed highest mortality (74%) with Agro NPK fertilizer.

Histopathology of the liver of fish exposed to Agro NPK fertilizer after 96hrs revealed severe damage and necrosis of the liver cells when compared with the control which had a normal distribution of hepatic cells. Similarly, results from histopathology of the gills of the exposed fish to Agro NPK fertilizer revealed fusion and damages to gills lamellae when compared with the control which has normal distribution of the gills lamellae. Severe changes in the epithelia of the gill arch were also observed. This finding corroborates with that of Govind, (2014) and Ekanem *et al.*, (2012). Findings from histopathology of the intestine revealed severe damages to the intestinal walls and mucosa when compared with the control fish which had a normal appearance as shown in plates 19-22. This finding agrees with the earlier reports of Ekanem *et al.*, (2012).

## Conclusion

Histopathology of the liver of the fish exposed to Agro NPK fertilizer revealed severe necrosis of the liver cells when viewed with an Amscope electron microscope at x10 magnification. Images from the control showed normal distribution of hepatic cells. Images from the histopathology of the gills revealed severe damages to the gill lamellae by the Agro NPK fertilizer whereas the control showed a normal distribution of gills lamellae. And in the same

vein, histopathology result on the effect of Agro NPK fertilizer on the intestine of the exposed fish revealed chronic damages to the intestinal mucosa when viewed with AMSCOPE electron microscope at x10 magnification. In all, a two way Analysis of variance on effect of Agro NPK fertilizer toxicant with different concentrations on histopathology of the catfishes at the end of 96hr bioassay showed significantly different at  $P < 0.05$  among the various concentrations studied.

**Conflict:**

The authors have declared that is no conflict of interest in this manuscript.

## References

Abbas, H. H. H. & Mahmood, H. M. (2003). The toxicological effect of water pollution on the Nile Tilapia fish (*Oreochromis niloticus* and *Clarias gariepinus*) exposed to mixtures of copper and salts. *Egyptian Journal of Basic & Applied Physiology*, 3, 89-106.

Abel, P. D. (2006). *Toxicity of synthetic Agro NPK fertilizer to fish and aquatic invertebrates*, 6(3), 279-298.

Adams, S. M., Shephard K. L., Greeley, M. S., Jr. Jimenez, B. D., Ryon, M. G., Shugart, L. R. & McCarthy, J. F. (1989). The use of bioindicators for assessing the effects of pollutant stress on fish. *Marine Environmental Research*, 28, 459-464.

Adewaye (2010). Comparative study on the behavioural responses of *Clarias gariepinus* fingerlings on exposure to soaps and Agro NPK fertilizer effluents. 1, 89-95.

Adewoye, and Lateef, A. (2004). Evaluation of the microbiological characteristics of Oyon river - A polluted river in North-central. *Pollution Research*, 23 (14), 587-591.

Adewoye. S. O., Fawole, O. O., Owolabi, O. D. & Omotosho, J. S. (2005). Toxicity of cassava wastewater effluents to African catfish: *Clarias gariepinus*. *Ethiopian Journal of Science*, 28 (7), 189 - 194.

Adham, K. G., Hamed, S. S., Ibrahim, H. M. & Saleh, R. A. (2002). Impaired functions of Nile Tilapia, *Oreochromis niloticus* from polluted waters. *Ada Hydromedica et Hydrobiologica*.

Aguiwo, J. N. (2002). The toxic effects of cymbush pesticides on growth and survival of African catfish, *Clarias gariepinus* (Bachel, 1822). *Journal of Aquatic Science*, 77(2), 81-84.

Akhtar, M. H. (1986): The deposition of  $^{14}\text{C}$  agrochemical residues in plants and animals tissue when consumed by animals. In T. Greenhalgh and T. R. Roberts (Eds.), *Pesticide Science and Biotechnology*. Ottawa: Proceedings Chemistry, 10 - 15.

Anastacio, P. M, Holten-Lutzhoft, H. C., Hailing-Sorensen, B. & Marques, J. C. (2000). Surfactant (Genapol OX-80), toxicity to *Selenastrum capricornulum*. *Chemosphere*, 40, 835-838.

Anderson, T., Forlin, L., Hardig, J. & Larsson, A. (1988). Physiological disturbances in fish living in coastal water polluted with bleached kraft pulp mill effluents. *Canadian Journal of Fisheries Aquatic Science*, 45, 1525-1536.

Antonio, F. F., Jorge, V. F. C., Sofia, G. S., Sandra M. M., Joao, C., Pedro, M. & Antonio, F. F. (2007). Histopathological changes in liver and gills epithelium of Nile tilapia, *Oreochromis niloticus*, exposed to waterborne copper. *Pesq. Vet.Bras.Marco*. 27(3), 103-109.

Avoajah, D. A. & Oti, E. E. (1997). Effect of sub-lethal concentration of some pesticides on the growth and survival of the fingerlings of the African freshwater catfish: Heteroclarias (Hybrid fingerlings) *Nigeria Journal of Biotechnology*, 8, 40-45.

Ayoola, S, O. (2008). Histopathological effects of glyphosate on Juvenile African catfish (*Clarias gariepinus*) *Ameri-surasian Journal of Agricultural and Environmental Science*, 4 (3), 362-367.

Abdel- Moneim A. M. (2005). Froese (2014). Behaviour, Habitat and Characteristics of the African catfish (*Clarias gariepinus*). 2<sup>nd</sup> edition, pages 3-5

Abel, P.D. (2006). Toxicity of synthetic Agro NPK fertilizer to fish and aquatic invertebrates, 6(3), 279-298

Adams, S. M. Shephard K. L., Greeley, M. S. Jr. Jimenez, B. D. Ryon, M. G. Shugart, L. R. McCarth, J. F. (1989). The use of bioindicators for assessing the effects of pollutant stress on fish. *Marine Environmental Research*. 28, 459-464.

Adewaye, A.O. (2010). Comparative study on the behavioural responses of *Clarias gariepinus* fingerlings on exposure to soaps and detergent effluents. (1) 89-95

Adewoye, and Lateef (2004) Evaluation of the microbial characteristics of the Oyun River- A polluted river in North-Central Nigeria. *Pollution Research*. 23(14), 587-591

Adewoye, S. O. and Favwole, O. O. (2002). Acute toxicity of soap and detergent effluent to fresh water. *Clarias gariepinus* fingerlings. *African J. Sc.* In press). 20(5): 22-35

Adewoye, S. O. Fawole, O. O. Owolabi, O. D and Omotosho, J. S. (2005). Toxicity of Cassava wastewater effluents to African catfish (*Clarias gariepinus*) *Ethiopian Journal of Science*, 28 (7), 189-194.

Adham, K. O. Hamed, S. Ibrahim, H.M. and Saleh, R. A. (2002). Impacted functions of Nile Tilapia, *Oreochromis niloticus* from polluted waters. *Acta Hydrochemica et Hydrobiologica*. Indian Journal of aquaculture 16 (2). 77-80

Aghoghovwia, O. A. (2015). Toxicity of detergent to the Aquatic environment. First edition, Nixon publishers, pages 12-13

Aguiwo, J. N. (2002). The toxic effects of cymbush pesticides on growth and survival of African catfish, *Clarias gariepinus* (Behell 1822). *Journal of Aquatic Science*, 17 (2), 81-84.

Andem, A. B, Odey, C. O, Esenowo Ime and Inyang Inimfon (2019). Ecological approach of plankton responses to water quality variables of a tropical River, South-Eastern Nigeria: A Bioindicator base of community Assessment of Idundu River. *Asian Journal of fisheries and aquaculture*. 3(1),1-17, 2019

Bucher, P. & Hofer, R, (1 994). The effects of treated domestics sewage on three organs (gills, kidney, liver) of brown trout (*Salmo irutta*). *Water Research*, 27, 255-261.

Byne, P., Speare, D. & Ferguson, H. W. (1989). Effects of cationic Agro NPK fertilizer on the gills and blood chemistry of rainbow trout. *Salmo Gaurdner*, 6, 185-196.

Canli, M. & Kargin, F. (1995). A comparative study on heavy metals (Cu, Cr, Pb and Ni) accumulates in the tissue of Carp. *Cyprinus carpio* and Nile fish. *Tilapia niloticus*. *Tropical Journal of Zoology*, 19, 165-171.

Boughlenger (1911), David 1935 Turgel, 1983 and Hogedoorn, 1980. A textbook on Characterization of the African Catfish, pages 25-36

Capkinset *al.*, (2009). Study on the sub chronic toxicity of composite inorganic fertilizers such as Ammonium sulfate on the skin, liver, gills, pancreas, kidney of Juvenile Rainbow trout fish (*onchorhynchus mykiss*).

Dambo, (1999). Effects of exposing aquatic organisms to crude and refined water soluble oils.

Ekanem and Oscar (2004). Effects of Agro NPK fertilizer on the behavioural responses of *Clarias gariepinus* sub adult in Calabar, Nigeria. *Asian Journal of Fisheries and Aquaculture*. (2) 4, 23-26

Ekanem, A. P and Ekpo, A. L., (2006). Effects of commercial detergent on the juveniles of *Heterbranchus longifilis* (Cuvier and Valenciennes). *African Journal of Environmental Pollution and Health*, 6 (1), 18-23.

Ekanem, A. P. (2012). Anatomy and physiology of the African catfish (*Clarias gariepinus*) Adults. *African Journal of Marine sciences* . 8 (1), 10-23.

Environmental protection Agency (2011). Chemical limits and standards in the aquatic ecosystem. Page 45-60

Esenowo, I. K. and Ugwumba, D. A. (2010) Growth response of catfish (*Clarias gariepinus*) exposed to water soluble fraction of detergent and diesel oil, *Environment Research Journal*, 4 (4), 298-301.

Ewa-Oboho, O. Amu (2009). Effectis of crude oil extracts on early stages of African catfish *heterobranchuslongifilis* (VAL) reared under controoled condition. Global journal on environmental sciences. 8(2), page 30-41

Farkas, A. Salanki, J. and Speeziar, (2007). Relation between growth and the heavy metals concentration in organs of bream, *Abramisbrama L.*, populating late Blaton. *Archive of Environmental Contaminants and Toxicology*.43 (2), 236- 243.

Francis E, A, Ita-EwaOboho Paul J. Udo and Ekaete F. Asuquo. Fish species used as biomakers for heavy metals and hydrocarbon contamination for Cross River State, Nigeria. *Environmentalist*.8(2). 30-41

Geetanjali, Choudhavy and Iha (2013). Influence of household Agro NPK fertilizer on some serum biochemical parameters of fresh water fish, chanapuntatus. *Asian journal of Biotechnology*. 21 (3), 121-123

George, R.O. (2012)Acute toxicity effects of Quo Iboe Crude on the gills of *Clariasgariepinus* juveniles. *Asian Journal of Biology* 1(3), page 34-40

Govind, P. O ( 2014). Effects of Linear Alkyl Benzene Sulfonate on the gills structure of the African catfish ( *Clariasgariepinus*). *Environment Research Journal*, 4 (4), 288-301.

Health 1990, Benneth (1990) Hematological and histopathological changes of *C. gariepinus*adults exposed to petroleum product. *Journal for environment andtoxicology*, 5(4), 20-25

Holzloenher, (2002). A textbook on wetlands and coastal zones in Calabar, First editon, page 40-45

Isyaku, B. and Solomon J. R., (2016).Effects of Agro NPK fertilizer and the growth of the African catfish *Clariasgariepinus*).

Ivon and Ndome, (2016).Comparative effects of Agro NPK fertilizer and liquiod soaps on the African catfish (*Clariasgariepinus*) fingerlings in Calabar, Nigeria.*Asian journal of Biology* 2(4), 12-18

Lenntech, (2012).*Bulletins detergent occuring in fresh water*.(vol. 6) Rotterdamsewg: Lenntech BV, 123.

Morah, F. N. (2012) *Advanced organic chemistry*, 3<sup>rd</sup> Edition Pp. 158.

National Population Commission. (2005). *Statistical update on the Nigerian population in Nigeria*, 2016 edition, volume 20, 12-13

Ndome, C. Mowang, D. A. and Ayibaerai, T. T. (2013). Comparative acute toxicity of local Agro NPK fertilizer (Omo and Ariel) on fingerlings of the *Clarias gariepinus* *Heterbranchus longifillsc hybrid*, *AACL Bioflux*, 6 (4), 413 - 420.

Ogundele, O., Ihuahi, J. A., Omojowo, F. S. Bitrus, P. (2005). Toxicity of linear *alkybenenesulphonate*(LAS) detergent, to *Clariagariepinus fingerlings*. In 19<sup>th</sup> Annual Conference of the Fisheries Society of Nigeria (FISON), 29<sup>th</sup> Nov. 03Dec. 2004, Ilorin, Nigeria, pp 271-276.

Ogundiran, M. A. Fawole, G. O. Adewoye, S. O. and Ayandiran, T. A. (2010). Toxicological impact of detergent effluent on juvenile of African catfish (*Clariagariepinus*) (Bushell 1822), *Agriculture and Biology Journal of North America 1* (3), 330 -342.

Ogundiran, M. A. Fawole, O. O. Adewoye S. O. & Ayandiran, T. A. (2009). Pathologic lesions in the gills of *Clariagariepinus* exposed to sub-lethal concentrations of soaps and detergent effluent. *Journal on cell and Animal Biology*, 3 (5), 078 -082.

Ogundiran, M. A. Fawole, O. O. and Adewoye, S. O. (2007). Effects of soap and detergent effluents on the haematological profiles of *Clarias gariepinus*, *Science Focus*, 12 (1), 84-88.

Omoregie, F. Ufodike, F. B. C. and Keke I. R. (1990). Tissues chemistry of *O. niloticus* exposed to sub lethal concentration of gammalin 20 and actellic 251EC. *Journal of Aquatic Science*, 5, 33-36.

Preeti B, Sexena M., (2016). Evaluation of lethal concentration of fertilizer (NPK) to fresh water fish, *mystus vitatus*. River State Environmental Protection Agency, (1992). Animal Environment reports.

Sangeetha, D.F(2011). Acute Toxicity of ammonium sulphate and composite fertilizer NKP 15:15:15 on the African catfish fingerlings.

Udejaja, O. Goodness, (2016). Effect of petroleum by products (Kerosene and fuel) on the hematology of African catfish (*Clariagariepinus*). *The Environmentalist*, 2(15), 25-36