



**A REVIEW ON THE PHYTOCHEMICAL,
PHARMACOLOGICAL, AND ANALYTICAL STUDIES CONDUCTED
ON *SEDUM LINEARE THUNB***

Sunil Kumar^{*1}, A.K.S. Rawat², Akash Ved³, Peeyush Bhardwaj⁴, Gyan Singh⁵

^{1,2}Maharishi School of Pharmaceutical Science, Maharishi University of Information
Technology, Sitapur road, Lucknow- 226013, Uttar Pradesh, India.

³Faculty of Pharmacy, Dr. APJ Abdul Kalam Technical University, Lucknow -226028, Uttar
Pradesh, India.

⁴Institute of Pharmacy, Bundelkhand University, Jhansi-284128, Uttar Pradesh, India.

⁵P. K. University, Shivpuri Road, Thanra- Shivpuri-473660, Madhya Pradesh, India.

Corresponding author

Mr. Sunil Kumar

PhD. Research Scholar,

Maharishi School of Pharmaceutical Science,

Maharishi University of Information Technology,

Sitapur road, Lucknow- 226013, Uttar Pradesh, India.

Email-sunilsinghpharmacy@gmail.com

ABSTRACT:

Within the context of traditional Chinese medicine, the use of the whole *Sedum lineare* Thunb. plant as a therapeutic agent is referred as "*Sedi linearis* Herba." This term originated in China. It was often used as a treatment for a wide range of medical conditions, such as hepatitis, swelling of the neck, dysentery, dermatitis rhus, burns, scalds, traumatic bleeding, and a huge number of other conditions. It has been shown that in addition to being able to modify thyroid hormone, this plant also contains the traits of being antibacterial, anticancer, nephroprotective, analgesic, anti-inflammatory, and anti-diarrheal. The capacity of the plant to shield the kidneys from injury is what's meant by the term "nephroprotective." Active chemical substances such as hyperoside, isoquercetin, astragalin, beta-amyrone, and oleanene triterpenes from *Sedum lineare* Thunb. have been identified from the plant, amongst a great many more. These compounds are what give the plant its medical qualities, and they are accountable for those qualities. In this review, we will be concentrating on the medicinal properties, pharmacological and analytical activities, and phytochemical make-up of *Sedum lineare* Thunb.

Keywords: *Sedum lineare* Thunb., Phytochemistry, δ -Amyrone, Pharmacological activity, Analytical study.

INTRODUCTION:

Traditional Chinese medicine using the whole *Sedum lineare* Thunb. plant is known as Sedi linearis Herba in China¹. It was often used as treatment for a wide range of medical issues, such as hepatitis, swelling of the throat, dysentery, dermatitis rhus, burns, scalds, traumatic bleeding, and so on^{2, 3}. The potential for *Sedi linearis* Herba to stop the formation of malignant cells has piqued the curiosity of the Tujia minority area in western Hubei, China⁴⁻⁶. Pharmacological research on mice with experimental acute liver injury found that *Sedi linearis* Herba protected the liver, reduced alanine amino transferase (ALT) activity, and cleared up their jaundice⁷⁻¹⁰. These results were seen in mice. The content of malondialdehyde (MDA) in the blood and the hepatic tissue was also dramatically reduced, and the activity of the superoxide dismutase enzyme was enhanced^{11, 12}. Furthermore, ethyl acetate extracted from the plant was shown to have a significant anticancer effect on many different types of cancer cells¹³⁻¹⁴. As an added bonus, the *Sedi linearis* Herb has a long history of use as a vegetable in some regions of China¹⁵. As a result, not only did this drug have a wide variety of therapeutic characteristics, but it was also safe to use in its crude medicinal form, arguing that it warranted further investigation and development¹⁶⁻¹⁸.

It's generally known that crude drugs include a wide variety of ingredients. Therefore, traditional quality control approaches, which primarily concentrate on quantifying a single active component, cannot provide a comprehensive evaluation of the raw drug's quality^{19, 20}.



FIGURE 1: PLANTS OF *SEDUM LINEARE THUNB.*²¹

Culture

Plants need full sunshine and soil that is medium to dry in order to flourish. Ideal for conditions with a considerable quantity of shade²². It can withstand very high temperatures as well as arid situations with ease²³. Sandier-than-gravel soils with a fertility level between medium and low are optimal for plant development²⁴. If you want to get satisfactory results, you will need to ensure that the soil drains well enough²⁵⁻²⁶. Allowing plants to flourish in their natural habitat may provide a beautiful ground cover²⁷. Locations in the St. Louis area, the northernmost latitude at which this plant may survive the winter, should be chosen with care due to the need for protection²⁸. This plant is hardy only in warm climates that never drop below freezing^{29, 30}.

Noteworthy characteristics

A mat-forming, evergreen stonecrop native to eastern Asia and known by its scientific name, *Sedum lineare* Thunb. It may reach a height of 4–6 inches and a width of 12 inches or more, depending on whether it develops stems that stay erect or fall over. Small, straight or lanceolate (up to 1 1/4 inches long) leaves "long) have a very light green colour. Flowering yellow plants (to 14.5 mm in "bloom cynically from late spring to early summer³¹⁻³³.

Most sedums grow in a sitting posture, therefore the name "genus" was derived from the Latin word for "to sit," "sedeo" (they sit and sprawl over rocks)³⁴.

The linear form of the leaves is reflected in the choice of the specific epithet³⁵.

The common name "stonecrop" is derived from the fact that many sedum species are found in the wild clinging to the sides of steep cliffs³⁶.

DISTRIBUTION:

Carpet Sedum, an evergreen succulent that thrives in full sun with no care, may be used in places where other plants die³⁷. The Latin words *sedeo*, meaning "to sit," and *lineare*, meaning "linear," combine to form the plant's name. Its native range includes parts of eastern Asia. Sedums are more often known as stonecrops³⁸. As a result, sedums are often found in their natural habitats on stony or rocky terrain. This tough plant can hold its own in scorching temperatures and dry conditions, and it has the potential to spread out in a low mat-like fashion³⁹⁻⁴¹. Its versatility makes it a great choice for a wide variety of planting situations, including shady regions, narrow spaces, shady corners, buried in or flowing over walls, rock gardens, xeriscapes, and hanging baskets⁴². As long as there is enough drainage, the Carpet Sedum may flourish in almost any soil type⁴³. In order to thrive, the soil should be sandy or gravelly. Some shade is OK, but the plant will grow more open and leggier as a consequence^{44, 45}.

Carpet Sedum is a plant that has been proposed as a viable option for green roofs on flat-roofed buildings in Shanghai, China. Carpet sedum has a shallow root system and hence cannot establish itself in a soilless environment⁴⁶.

The Carpet Sedum is easily reproduced throughout the year by layering, stem cutting, and division⁴⁷. When a leaf is placed on top of dirt, roots develop at the point of contact. The leaf will stay there if you sprinkle a little dirt on top of it⁴⁸. You may make your plants larger by dividing them and replanting the cuttings. It is possible to do a stem cut by simply taking off a part of the plant and replanting it⁴⁹. Carpet Sedum may spread naturally from plant to plant by means of its own seeds⁵⁰. The Carpet Sedum is not only pest-proof but also blooms a little yellow flower from late spring into early summer. Pollinators are attracted to this flower⁵¹. It is recommended to keep a watch out for slugs, snails, and scale, although these insects are not generally considered to be a major problem⁵².

Origin and Habitat: The countries of Japan, China (including the provinces of Anhui, Fujian, Gansu, Guangdong, Guizhou, Henan, Hubei, Hunan, Jiangsu, Jiangxi, Shaanxi, Sichuan, and Yunnan), and Eastern Asia⁵³. The plant was brought to and eventually naturalised in Eastern Europe and the United States (Georgia). This species was first

documented using Japanese specimens, but its origins are unknown. The species was described, however, based on data collected in Japan⁵⁴.


Cultivation and Propagation: *Sedum linearare* Thunb, a kind of moss-like ground cover, is an excellent option for use in rock gardens, stepping-stone containers, and wall niches. Landscape elements that serve as a border, such as rocks or a garden⁵⁵. Use them in bulk or in clusters for the best results. Great as a ground-based scrambler, especially when taught to traverse or circle a nearby rock, so long as the dirt is well drained. Its stems may grow to a length of roughly 30 centimetres, making it a superb plant to use in a hanging basket. Sedums are a kind of succulent that is easy to grow and tolerates a broad range of environmental factors⁵⁶. They do well in full sun or partial shade, wet or dry soil, and many intensities of light. However, they only blossom into their full splendour when given sufficient water and sunshine, and they thrive best when cultivated in the fresh air⁵⁷⁻⁶⁰. Due to its ability to withstand cold and drought, minimal soil requirements, and lack of penetrating roots, this plant has been proposed as a good option for "greening" Shanghai, China's flat-roofed skyscrapers⁶¹⁻⁶².

MEDICINAL USE OF *SEDUM LINEARE THUNB*:

A fresh plant may be crushed and used to burns and scalds for relief⁶³. *Sedum linearare* Thunb. a common garden plant, was used historically to treat a broad range of inflammatory skin diseases. Many of the leaves' constituents are used to speed wound recovery and calm irritated skin. Before using the leaves, their protective cuticle is stripped away⁶⁴.

PHYTOCHEMISTRY OF *SEDUM LINEARE THUNB*:


Through the use of phytochemical analysis, it was shown that *Sedum linearare* Thunb. plants contained hyperoside, isoquercetin, and astragalin. In addition to two known oleanene triterpenes, -amyrone (2) and -amyrine acetate, an alcoholic extract of the entire plant of *Sedum linearare* Thunb revealed the presence of one novel olean-13(18)-ene-3,12,19-trione (1). This discovery was made possible by the presence of an additional oleanene triterpene. These findings came from the alcoholic extract's petroleum ether fraction, which was separated out and analysed (3)⁶⁵⁻⁶⁷.




Hyperoside




Isoquercetin



Astragalin



δ -Amyrone



Triterpene oleanenes derived from *Sedum lineare* Thunb.

PHARMACOLOGICAL AND ANALYTICAL PROPERTIES OF *SEDUM LINEARE* THUNB:

For this chromatography, we used a BDS Hypersil C18 column and a mobile phase of acetonitrile-0.1% acetic acid solution (gradient elution). The results revealed substantial differences in the concentrations of the major active component across samples collected from the same environment throughout each of the four seasons. Samples obtained towards the end of April had a substantially greater ingredient content than those tested in the middle of August. Furthermore, the major active component concentrations varied across samples collected during the same season but from different locations. The sample's concentrations of the most significant elements, for example, were greater in August's rainy climate than in the dry environment. These results, taken together, provided conclusive evidence that the HPLC fingerprint analysis and contents determination technique that had been devised was helpful for analysing and maintaining control over the quality of *Sedum lineare* Herba⁶⁸.

Following the same detection method and a number of collection seasons, HPLC fingerprints of *Sedum lineare* and *Sedum sarmentosum* were produced and compared. *Sedum lineare* has been shown to contain hyperin, isoquercitrin, and astragaloside, all of which were detected in an HPLC analysis of the plant's fingerprint. While fingerprints from *Sedum lineare* grown in the same location but in different environments were nearly indistinguishable, fingerprints from the same plant grown at different times of the year showed striking differences, with the area of the most common peaks shifting from small to large and the number of peaks changing slightly. Despite the fact that they shared four peaks in their HPLC fingerprint, the two *Sedum* species could be clearly distinguished from one another. These two kinds are best

harvested during full bloom. The established method has promise as a standard for recognising and evaluating *Sedum lineare* Thunb⁶⁹.

The whole *Sedum lineare* Thunb plant has been utilised as a traditional folk medicine for the treatment of seventeen various conditions, including sore throat, recurring hepatitis, jaundice, and dysentery. In 19 different animal models of inflammation, the pentacyclic triterpene molecule -amyrone (13(18)-oleanen-3-one) isolated from *S. lineare* Thunb was shown to have a potent anti-inflammatory effect. Twenty mice had decreased xylene-induced ear edoema when pretreatment with -Amyrone (intraperitoneally), -amyrone decreased levels of nitric oxide (NO), prostaglandin E2 (PGE2), 21 interleukin-6 (IL-6), and leukocyte counts in acetic acid-induced peritonitis in vivo. The purpose of this study was to investigate the effect of - Amyrone in lipopolysaccharide (LPS)-generated perito- 23 neal macrophages and to determine its possible 22 mechanism. The data indicated that -Amyrone effectively suppressed the production of IL-6, TNF-, and NO. In addition, the results revealed that protein-level regulation of cyclooxygenase-2 is mediated by -amyrone rather than cyclooxygenase-1 (COX-1) (COX-2). Results like these suggest that -amyrone is a bioactive molecule with anti-inflammatory effects that may contribute to COX-2 regulation⁷⁰.

Two known oleanene triterpenes, -amyrone (2) and -amyrine acetate (3), and one new olean-13(18)-ene-3,12,19-trione (1) were isolated from a petroleum ether fraction of an alcoholic extract of the whole plant of *Sedum linear* Thunb (3). X-ray diffraction analysis was used to verify the structure of the unique molecule, and further spectroscopic methods, such as 1D and 2D NMR and HR-ESI-MS, provided more insight into the molecule's composition. NMR analysis allowed for the identification of the proven ones by comparing the compounds' profiles to those of already-identified chemicals. The compounds' anti-TNF- and anti-NO effects were also evaluated *in-vitro*⁷¹.

Important for healthy plant growth, zinc (Zn) might have the opposite effect depending on the concentration. The morphological characteristics and antioxidant capabilities of *Sedum lineare* Thunb. were investigated at varying concentrations of zinc in relation to 24-Epibrassinolide (EBR). 24-Epibrassinolide (EBR) is a well-known steroid phytohormone that regulates plant development and alleviates the damage caused by abiotic stress. Several morphological features were significantly improved by simultaneous foliar treatment with 0.75 M EBR compared to plants exposed to Zn alone, and the benefits of this therapy on plant growth were most pronounced at high Zn concentrations. Increases of 111%, 85%, and 157% in stem length, fresh weight, and internode length were seen in EBR-treated plants compared to Zn alone-treated plants at a toxic 800 M Zn. EBR spray reduced the accumulation of reactive oxygen species (ROS) and lipid peroxidation in plants. In addition, it increased the activities of antioxidant enzymes such as peroxidase (POD), ascorbate peroxidase (APX), and glutathione reductase (GR), as well as the contents of antioxidative agents such as ascorbic acid (AsA) and glutathione. These results suggest that EBR spray may be a useful tool for the (GSH). By demonstrating that EBR helped *S. lineare* Thunb. resist high-zinc stress by strengthening the antioxidant system, the results of this research

provided novel insight into how to increase the genetic diversity of Crassulaceae plants cultivated in heavy metal-contaminated soil for phytoremediation. This research was funded by the National Science Foundation⁷².

We utilised acetonitrile as the stationary phase and a 0.1% acetic acid solution in acetonitrile as the mobile phase (gradient elution). The results revealed substantial differences in the concentrations of the major active component across samples collected from the same environment throughout each of the four seasons. Samples obtained towards the end of April had a substantially greater ingredient content than those tested in the middle of August. Furthermore, the major active component concentrations varied across samples collected during the same season but from different locations. The sample's concentrations of the most significant elements, for example, were greater in August's rainy climate than in the dry environment. Collectively, these findings proved that the HPLC fingerprint analysis and contents determination approach established was useful for analysing and controlling the quality of *Sedi linearis* Herba⁷³.

Recent advances notwithstanding, neuropathic pain continues to be a pressing issue in global health. As a result, it is crucial to look into other methods of relieving neuropathic pain. The role of neuroinflammation in the development of neuropathic pain has been well acknowledged. The Chinese herb *Sedum lineare* Thunb. (SLT), which comes from the whole grass of a Crassulaceae plant, is said to have anti-inflammatory effects. However, it is not known how SLT may help with neuropathic pain. In this study, neuropathic pain was induced in rats using spared nerve injury (SNI).

In this study, SLT (p.o.) was administered once a day to SNI rats for a total of 14 days. Pain-related behaviours in the animals were measured using the paw withdrawal threshold (PWT) and the CatWalk gait characteristics. A Western blotting test was used to determine the levels of inflammatory mediators and pain-related signalling molecules that are expressed in the spinal cord. Mechanical hypersensitivity in SNI rats was decreased by SLT treatment in a dose-dependent manner. Doses of SLT (30, 100, and 300 mg/kg, p.o.) were effective. The optimum dose of SLT, 100 mg/kg, successfully suppressed microglial activation, lowered spinal HMGB1, TLR4, MyD88, TRAF6, IL-1, IL-6, and TNF-protein levels, and improved gait parameters in SNI rats.

Protein expression for IB and IL-10 was unquestionably upregulated after SLT treatment, whereas phosphorylation levels for IKK and NF-kappaB p65 were markedly lowered in the spinal cord of SNI rats. Combined, our findings suggest that SLT may work as a therapeutic drug for neuropathic pain by inhibiting the spinal TLR4/NF-B signalling pathway in SNI rats⁷⁴.

CONCLUSION:

There has been a significant amount of research conducted on *Sedum lineare* (Thunb.), all of which indicates that it has a massive amount of untapped biological potential. There is reason to be hopeful that the evidence for the use of this plant in medicine may be located in the particulars that are given in this analysis of the phytochemical and other biological

characteristics of the extracts. Geographical areas have a role in both the diversity of phytochemical compounds that *Sedum lineare* Thunb has and the efficacy of the therapeutic characteristics that it possesses.

Even in this day and age, the bulk of the world's population relies almost exclusively on plant-based medicines as their sole supply of pharmaceuticals. Because of this, it is still challenging for scientists to deliver medication that is not only inexpensive but also effective and safe, especially for those who live in rural regions. More study has to be done on these species of *Sedum* as well as their quantification of individual phytoconstituents and their pharmacological profiles, which are based on *in-vitro* research, *in-vivo* research, and clinical trials.

REFERENCES

1. Liu D, Mei Q, Long W, Wan X, Wan D, Wang L. HPLC Fingerprint Analysis and Content Determination of Extract with Anticancer Activities of *Sedi Linearis* Herba. *Pharmacognosy Journal*. 2017;9(2).
2. Wang KR, Ding R, Lin J, Hu DJ, Zhong SH. A cross-national comparison of medicinal plants used by the Miao, Yi and Lisu ethnic groups in Yanbian, China.
3. Li M, Wang A, Zhang Y, Han T, Guan L, Fan D, Liu J, Xu Y. A comprehensive review on ethnobotanical, phytochemical and pharmacological aspects of *Rhus chinensis* Mill. *Journal of Ethnopharmacology*. 2022 Jul 15;293:115288.
4. De Philippis R, Vincenzin M. New and traditional energy resources from microbial activities in the agroindustrial system. *Italian Journal of Agronomy*. 2009 Apr 9;4(s1):141-6.
5. De Philippis R. Idrogeno e metano di originebiologica: microrganismi e processibiotecnologici per il recupero di energiadairesiduivegetali. *Idrogeno e metano di originebiologica: microrganismi e processibiotecnologici per il recupero di energiadairesiduivegetali*. 2014:7-21.
6. Patriarca C, Miritana VM, Gorrasi S, Fenice M. Trattamentobiologico di biomasselignocellulosiche e chitinose per la produzione di idrogeno e/o metano. *Ricerca di Sistema Elettrico ENEA, RapportotecnicoRdS/PAR2013/252*. 2014 Sep.
7. Meng X, Qiu Q, Zhen HS, Huang F, Zhen D, Jiang LJ, Ye M, Liu Y. Fingerprint and multi-index content determination of ethyl acetate extract of *Sedum emarginatum*Migo. *Pakistan Journal of Pharmaceutical Sciences*. 2021 Sep 3;34.
8. Liu LJ, Liu JN. A strategy for quality control of *Menispermum dauricum* DC based on cytotoxic activity and HPLC fingerprint analysis. *Indian Journal of Pharmaceutical Sciences*. 2016 Jan;78(1):143.
9. Yucui MA, Chun WA, Wei WA, Cui WU, Ning DU, Wen SU, Hui LI, Zhimao CH. An HPLC Fingerprint Identification of Dried Barks of *Ilex rotunda* and *Ilex godajam*. *Medicinal Plant*. 2018 Jun 1;9(3).
10. Hernadi E, Rohaeti E, Rafi M, Wahyuni WT, Putri SP, Fukusaki E. HPLC fingerprinting coupled with linear discriminant analysis for the detection of

- adulteration in *Orthosiphon aristatus*. Journal of Liquid Chromatography & Related Technologies. 2019 Oct 2;42(15-16):513-20.
11. Abdel-Hamid M, Osman A, El-Hadary A, Romeih E, Sitohy M, Li L. Hepatoprotective action of papain-hydrolyzed buffalo milk protein on carbon tetrachloride oxidative stressed albino rats. Journal of dairy science. 2020 Feb 1;103(2):1884-93.
 12. Goodarzi Z, Karami E, Yousefi S, Dehdashti A, Bandegi AR, Ghanbari A. Hepatoprotective effect of atorvastatin on Cadmium chloride induced hepatotoxicity in rats. Life sciences. 2020 Aug 1;254:117770.
 13. Abraham NN, Kanthimathi MS, Abdul-Aziz A. Piper betle shows antioxidant activities, inhibits MCF-7 cell proliferation and increases activities of catalase and superoxide dismutase. BMC complementary and alternative medicine. 2012 Dec;12(1):1-1.
 14. Poofery J, Khaw-On P, Subhawa S, Sripanidkulchai B, Tantraworasin A, Saeteng S, Siwachat S, Lertprasertsuke N, Banjerdpongchai R. Potential of Thai herbal extracts on lung cancer treatment by inducing apoptosis and synergizing chemotherapy. Molecules. 2020 Jan 6;25(1):231.
 15. Buhner SH. Herbal antibiotics: natural alternatives for treating drug-resistant bacteria. Storey Publishing; 2012 Jul 17.
 16. Živković J, Ilić M, Šavikin K, Zdunić G, Ilić A, Stojković D. Traditional use of medicinal plants in South-Eastern Serbia (Pčinja District): Ethnopharmacological investigation on the current status and comparison with half a century old data. Frontiers in Pharmacology. 2020 Jul 8;11:1020.
 17. Khan SA, Al-Balushi K. Combating COVID-19: The role of drug repurposing and medicinal plants. Journal of infection and public health. 2021 Apr 1;14(4):495-503.
 18. Stella B, Baratta F, Della Pepa C, Arpicco S, Gastaldi D, Dosio F. Cannabinoid formulations and delivery systems: Current and future options to treat pain. Drugs. 2021 Sep;81(13):1513-57.
 19. Schulze B, Jeon Y, Kaserzon S, Heffernan AL, Dewapriya P, O'Brien J, Ramos MJ, Gorji SG, Mueller JF, Thomas KV, Samanipour S. An assessment of quality assurance/quality control efforts in high resolution mass spectrometry non-target workflows for analysis of environmental samples. TrAC Trends in Analytical Chemistry. 2020 Dec 1;133:116063.
 20. Li Y, Shen Y, Yao CL, Guo DA. Quality assessment of herbal medicines based on chemical fingerprints combined with chemometrics approach: A review. Journal of Pharmaceutical and Biomedical Analysis. 2020 Jun 5;185:113215.
 21. https://species.wikimedia.org/wiki/Sedum_lineare.
 22. Zhang Y, Liao H. Epibrassinolide improves the growth performance of *Sedum lineare* upon Zn stress through boosting antioxidative capacities. Plos one. 2021 Sep 7;16(9):e0257172.

23. Wu XX, Chen JL, Xu S, Zhu ZW, Zha DS. Exogenous 24-epibrassinolide alleviates zinc-induced toxicity in eggplant (*Solanum melongena* L.) seedlings by regulating the glutathione-ascorbate-dependent detoxification pathway. *The Journal of Horticultural Science and Biotechnology*. 2016 Jul 3;91(4):412-20.
24. Alam P, Balawi TA. 24-Epibrassinolide (EBR) reduces oxidative stress damage induced by cadmium toxicity by restricting cd uptake and modulating some key antioxidant enzymes in maize plants. *Pakistan Journal of Botany*. 2021;53(1):56-66.
25. Jan S, Noman A, Kaya C, Ashraf M, Alyemeni MN, Ahmad P. 24-Epibrassinolide alleviates the injurious effects of Cr (VI) toxicity in tomato plants: Insights into growth, physio-biochemical attributes, antioxidant activity and regulation of Ascorbate–glutathione and Glyoxalase cycles. *Journal of Plant Growth Regulation*. 2020 Dec;39(4):1587-604.
26. Jan S, Alyemeni MN, Wijaya L, Alam P, Siddique KH, Ahmad P. Interactive effect of 24-epibrassinolide and silicon alleviates cadmium stress via the modulation of antioxidant defense and glyoxalase systems and macronutrient content in *Pisum sativum* L. seedlings. *BMC plant biology*. 2018 Dec;18(1):1-8.
27. Colares GS, Dell'Osbel N, Wiesel PG, Oliveira GA, Lemos PH, da Silva FP, Lutterbeck CA, Kist LT, Machado ÊL. Floating treatment wetlands: A review and bibliometric analysis. *Science of the Total Environment*. 2020 Apr 20;714:136776.
28. Ruthsatz B, Schitteck K, Backes B. The vegetation of cushion peatlands in the Argentine Andes and changes in their floristic composition across a latitudinal gradient from 39 S to 22 S. *Phytocoenologia*. 2020 Oct 5;50(3):249-78.
29. Fresco N, Bennett A, Bieniek P, Rosner C. Climate Change, Farming, and Gardening in Alaska: Cultivating Opportunities. *Sustainability*. 2021 Nov 17;13(22):12713.
30. Baxter LL, Anderson WF, Gates RN, Rios EF, Hancock DW. Moving warm- season forage bermudagrass (*Cynodon* spp.) into temperate regions of North America. *Grass and Forage Science*. 2022 Jun;77(2):141-50.
31. Najafloo R, Behyari M, Imani R, Nour S. A mini-review of Thymol incorporated materials: Applications in antibacterial wound dressing. *Journal of Drug Delivery Science and Technology*. 2020 Dec 1;60:101904.
32. Rajeshkumar S, Jeevitha M, Sheba D, Nagalingam M. Bacterial and fungal mediated synthesis, characterization and applications of AgNPs. In *Agri-Waste and Microbes for Production of Sustainable Nanomaterials* 2022 Jan 1 (pp. 165-186). Elsevier.
33. Rather RA, Bano H, Padder SA, Perveen K, Al Masoudi LM, Alam SS, Hong SH. Anthropogenic impacts on phytosociological features and soil microbial health of *Colchicum luteum* L. an endangered medicinal plant of North Western Himalaya. *Saudi Journal of Biological Sciences*. 2022 Apr 1;29(4):2856-66.
34. Blanusa T, Monteiro MM, Fantozzi F, Vysini E, Li Y, Cameron RW. Alternatives to Sedum on green roofs: Can broad leaf perennial plants offer better ‘cooling service’?. *Building and Environment*. 2013 Jan 1;59:99-106.

35. Trigas P, Kalpoutzakis E, Kalogiannis E, Valli AT, Kougioumoutzis K, Katopodis K, Constantinidis T. Noteworthy new floristic records from Greece. *Botanica Serbica*. 2021;45(2):321-31.
36. Cui X, Shen Y, Yang Q, Kawi S, He Z, Yang X, Wang CH. Simultaneous syngas and biochar production during heavy metal separation from Cd/Zn hyperaccumulator (*Sedum alfredii*) by gasification. *Chemical Engineering Journal*. 2018 Sep 1;347:543-51.
37. Frank G. *Striking Succulent Gardens: Plants and Plans for Designing Your Low-Maintenance Landscape [A Gardening Book]*. Ten Speed Press; 2021 Jan 26.
38. Doyle L. *Essential Green Roof Construction: The Complete Step-by-step Guide*. New Society Publishers; 2021 Nov 9.
39. Körner C. *Alpine plant life: functional plant ecology of high mountain ecosystems*. Springer Nature; 2021 Mar 31.
40. Dvorak B, Rottle ND. Green Roofs in Puget Lowland Ecoregions. In *Ecoregional Green Roofs 2021* (pp. 391-449). Springer, Cham.
41. Norris KD. *New Naturalism: Designing and Planting a Resilient, Ecologically Vibrant Home Garden*. Cool Springs Press; 2021 Jan 5.
42. Don M. *The Complete Gardener: A Practical, Imaginative Guide to Every Aspect of Gardening*. Dorling Kindersley Ltd; 2021 Mar 4.
43. Petrovich KE, Vladimirovna SM, Stolbovaya M, Kolpak E. The edges of the leaf blade. The main forms of the leaf blade. How it works. *Gas*. 2021 May 5.
44. Beecham S, Razzaghmanesh M, Bustami R, Ward J. The role of green roofs and living walls as WSUD approaches in a dry climate. In *Approaches to Water Sensitive Urban Design 2019* Jan 1 (pp. 409-430). Woodhead Publishing.
45. Heatherington C, Johnson A. *Habitat Creation in Garden Design: A guide to designing places for people and wildlife*. The Crowood Press; 2022 Sep 26.
46. Chatters C. *Flowers of the forest: Plants and people in the New Forest National Park*. Princeton University Press; 2021 Oct 12.
47. Jittin V, Tripti SR, Bahurudeen A, Hammadhu RJ. Cleaner construction of durable green rooftop in residential buildings with municipal water supply. *Cleaner Materials*. 2022 Sep 1;5:100125.
48. Ayilara MS, Olanrewaju OS, Babalola OO, Odeyemi O. Waste management through composting: Challenges and potentials. *Sustainability*. 2020 Jan;12(11):4456.
49. Smith M. *The Plant Propagator's Bible: A Step-by-Step Guide to Propagating Every Plant in Your Garden*. Cool Springs Press; 2021 Jun 29.
50. Nagase A, Dunnett N, Choi MS. Investigation of weed phenology in an establishing semi-extensive green roof. *Ecological engineering*. 2013 Sep 1;58:156-64.
51. Green K. *Plantiful: Start Small, Grow Big with 150 Plants that Spread, Self-sow, and Overwinter*. Timber Press; 2014 Jan 28.

52. Flint ML. Pests of the garden and small farm: A grower's guide to using less pesticide. UCANR Publications; 2018.
53. Fan JL, Wang JX, Hu JW, Wang Y, Zhang X. Optimization of China's provincial renewable energy installation plan for the 13th five-year plan based on renewable portfolio standards. *Applied Energy*. 2019 Nov 15;254:113757.
54. Montagnani C, Gentili R, Smith M, Guarino MF, Citterio S. The worldwide spread, success, and impact of ragweed (*Ambrosia* spp.). *Critical Reviews in Plant Sciences*. 2017 May 4;36(3):139-78.
55. Lu J, Yuan JG, Yang JZ, Chen AK, Yang ZY. Effect of substrate depth on initial growth and drought tolerance of *Sedum lineare* in extensive green roof system. *Ecological Engineering*. 2015 Jan 1;74:408-14.
56. Hernández-Zamora AK, Rodríguez-Dorantes A. Image Analysis of Root Induction in *Sedum praealtum* Cuttings. *European Journal of Biology and Biotechnology*. 2020 Nov 30;1(6).
57. Almusaed A. Biophilic and bioclimatic architecture: analytical therapy for the next generation of passive sustainable architecture. Springer Science & Business Media; 2010 Dec 21.
58. Van Jaarsveld E. Waterwise gardening in South Africa and Namibia. Penguin Random House South Africa; 2013 Oct 29.
59. Almusaed A. Introduction on Plants and Vegetations. In *Biophilic and Bioclimatic Architecture 2011* (pp. 47-83). Springer, London.
60. Rainer T, West C. Planting in a post-wild world: Designing plant communities for resilient landscapes. Timber Press; 2015 Oct 7.
61. McKenzie L. Design, Context and Use of Public Space: The Influence of Heat on Everyday Behaviour in Outdoor Settings—A Western Sydney Case Study. Unpublished PhD Thesis, University of New South Wales. 2017 Aug.
62. Kent JL, Thompson S. Planning Australia's healthy built environments. Routledge; 2019 Mar 18.
63. Yue-ling M, Yu-jie C, Ding-rong W, Ran X. HPLC determination of quercetin in three plant drugs from genus *Sedum* and conjecture of the best harvest time. *Pharmacognosy Journal*. 2017;9(6).
64. Enriquez-Ochoa D, Sánchez-Trasviña C, Hernández-Sedas B, Mayolo-Deloisa K, Zavala J, Rito-Palomares M, Valdez-García JE. Aqueous two-phase extraction of phenolic compounds from *Sedum dendroideum* with antioxidant activity and anti-proliferative properties against breast cancer cells. *Separation and Purification Technology*. 2020 Nov 15; 251:117341.
65. Tian L, SU J, Zhong C, Xie Y. Study on the Chemical Constituents of *Sedum lineare*. *China Pharmacy*. 2016:2956-8.
66. Niu XF, Liu X, Pan L, Qi L. Oleanene triterpenes from *Sedum lineare* Thunb. *Fitoterapia*. 2011 Oct 1;82(7):960-3.

67. Chiocchio I, Poli F, Governa P, Biagi M, Lianza M. Wound healing and in vitro antiradical activity of five *Sedum* species grown within two sites of community importance in Emilia-Romagna (Italy). *Plant Biosystems-An International Journal Dealing with all Aspects of Plant Biology*. 2019 Jul 4;153 (4):610-5.
68. Liu D, Mei Q, Long W, Wan X, Wan D, Wang L. HPLC Fingerprint Analysis and Content Determination of Extract with Anticancer Activities of *Sedi Linearis* Herba. *Pharmacognosy Journal*. 2017; 9(2).
69. Lu LQ, Mei Q, Wan DR, Yang XZ, Qiao S, Zhao YD. HPLC characteristic fingerprints of *sedi linearis herba* and *sediherba*. *Zhong yaocai= Zhongyaocai= Journal of Chinese Medicinal Materials*. 2014 Apr 1;37(4):583-7.
70. Niu X, et al, δ -Amyrone, a specific inhibitor of cyclooxygenase-2, exhibits anti-inflammatory effects in vitro and in vivo of mice, *Int Immunopharmacol* (2014), <http://dx.doi.org/10.1016/j.intimp.2014.04.019>.
71. Niu XF, Liu X, Pan L, Qi L. Oleanene triterpenes from *Sedum lineare* Thunb. *Fitoterapia*. 2011 Oct;82(7):960-3. doi: 10.1016/j.fitote.2011.05.011. Epub 2011 May 23. PMID: 21624444.
72. Zhang Y, Liao H (2021) Epibrassinolide improves the growth performance of *Sedum lineare* upon Zn stress through boosting antioxidative capacities. *PLoS ONE* 16(9): e0257172. <https://doi.org/10.1371/journal.pone.0257172>.
73. Liu D, Mei Q, Long W, Wan X, Wan D, Wang L. HPLC Fingerprint Analysis and Content Determination of Extract with Anticancer Activities of *Sedi Linearis* Herba. *Pharmacogn J*. 2017; 9(2):128-34.
74. Wang XY, Ma HJ, Xue M, Sun YL, Ren A, Li MQ, Huang ZH, Huang C. Antinociceptive effects of *Sedum lineare* Thunb. on spared nerve injury-induced neuropathic pain by inhibiting TLR4/NF- κ B signaling in the spinal cord in rats. *Biomedicine & Pharmacotherapy*. 2021 Mar 1; 135:111215.