



## ASSESSMENT OF VARIOUS RISK HABITS AMONG ORAL CANCER PATIENTS

Miloni Suresh Shah<sup>[a]</sup>, Mahathi Neralla<sup>[b]\*</sup>, Maragathavalli Gopal<sup>[c]</sup>

**Article History:** Received: 23.02.2022

Revised: 22.03.2022

Accepted: 21.04.2022

**Abstract:** Oral cancer is a major public health problem in the Indian subcontinent, where it ranks among the top three types of cancer in the country. Oral cancer is defined as the cancer of lips, mouth and tongue. The difference in incidence and pattern of oral cancer can be due to an overall effect of ageing of population as well as some regional differences in the prevalence of specific risk factor. Therefore the aim of the present study was to assess oral cancer incidence in patients based on forms of habits. This study was conducted in the Oral Cancer Department, Saveetha Dental College, Chennai. The study involved 50 patients who were clinically diagnosed with oral cancer. The various forms of habits observed were Guthka, Pan chewing, Smoking. Pearson chi square test was done to statistically analyze the data and to identify any significant level of association. The most predominant form of habit was observed to be Pan chewing (n=11;22%) Out of the total 50 patients involved in the study it was observed that it was most predominant in the males (78%) with Pan chewing being the most predominant habit (n=10;25.6%) More number of oral cancer with various forms of habits were observed in above 50 years of age group (n=29;58%) Within the limitations of our study it was observed that oral cancer incidence in patients based on forms of habits was observed in above 50 years of age with a high predominance in the males and Pan chewing being the most common form of habit. Further studies with a larger sample size are required to compare the efficacy of oral cancer based on various forms of habits.

**Keywords:** Cancer, Forms of habits, Prevalence, predominance, innovative study.

- [a]. <sup>1</sup>Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences (SIMATS), Saveetha university, Chennai- 77, India.  
 [b]. <sup>2</sup>Reader, Department of Oral and Maxillofacial Surgery, Saveetha dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences (SIMATS), Saveetha university, Chennai- 77, India.  
 [c]. <sup>3</sup>Head of department, Department of Oral Medicine and Radiology, Saveetha Dental college and Hospitals, Saveetha Institute of Medical and Technical Sciences (SIMATS), Saveetha university, Chennai-77, India.

**\*Corresponding Author**

**Email:** 151501038.sdc@saveetha.com, mahatin.sdc@saveetha.com, drgopalvalli@gmail.com

**DOI:** 10.31838/ecb/2022.11.02.005

### INTRODUCTION

Carcinoma is defined as the uncontrollable growth of cells that invade and cause damage to the surrounding tissues. Globally, oral carcinoma is the sixth most common cancer. Among the modern epidemics, oral cancer is the commonest cause of morbidity and mortality in developing countries <sup>1-2</sup>. Tobacco use and excessive alcohol consumption have been estimated to account for about 90% of cancer in the oral cavity, the oral malignancy risk increases when tobacco is used in combination with alcohol or areca nut <sup>3</sup>. The risk for developing oral malignancy is three times higher in smokers compared with non-smokers <sup>4</sup>. Major risk factors for oral cancer are the use of tobacco, betel quid, and alcohol <sup>5,6</sup>. Although existing tobacco and alcohol control policies, mouth cancer incidence has been increasing in most population-based cancer registries (PBCRs) in India. In a

country such as India, where access to health-care services and cancer-related awareness is highly variable, changes in incidence rates should be interpreted carefully <sup>7-8</sup>.

Oral cancers not only affect the cosmesis of the patients, but also the patient's lifestyle, life expectancy, patient's ability to communicate, major functions such as swallowing and chewing. In developing countries, the survival rate in patients with oral carcinoma is poor because of advanced clinical stages and advanced age at which the disease is diagnosed due to low socioeconomic status and awareness among the people <sup>9</sup>. Oral squamous cell carcinoma shows geographical variation with respect to the age, sex, site, and habits of the population <sup>10</sup>. In South East Asia, the high incidence was due to risk habits of smoking, betel quid, and tobacco chewing habits. The mixed habits act as a synergistic effect, with higher risk than independent risk habits. India had the highest incident rate of oral cancer; it was aptly labeled oral cancer capital of the world with an estimated 1% of the population having oral premalignant lesions <sup>11-12</sup>.

The most common habits related to oral cancer in India are use of tobacco smoking and tobacco chewing. Use of alcohol can be considered as an additive factor. As per the report obtained from the Global Adult Tobacco Survey, from Indian states and union territories conducted during 2009–2010, approximately 274.9 million people use tobacco in India. As per data, more than one-third (35%) of adult use tobacco in some form or the other (Smoked or smokeless), 163.7 million are users of only smokeless tobacco, 68.9 million are only smokers, and 42.3 million are users of both smoking and smokeless tobacco <sup>13,14</sup>. Early detection of these oral cancers will enable appropriate clinical management and monitoring. Moreover, improving the incidence, mortality, and survival rates of oral cancer requires a multi-tier structural approach that targets society, dentists, communities, and the individual. <sup>15-29</sup>

Even though clinical diagnosis occurs via examination of the oral cavity and tongue which is accessible by current

diagnostic tools, the majority of cases present to a healthcare facility at later stages of cancer subtypes, thereby reducing chances of survival due to delays in diagnosis<sup>30</sup>

Oral cancer will remain a major health problem and efforts towards early detection, and prevention will reduce this burden. This information gives us a knowledge of the burden of cancer, etiology studies and the effectiveness of the activities that have been undertaken to control cancer. Previously our team has a rich experience in working on various research projects across multiple disciplines<sup>31-40</sup> Now the growing trend in this area motivated us to pursue this project. The aim of this study is to assess the incidence of oral cancer based on various forms of habits.

## MATERIALS AND METHODS

**Study design and setting:** This is a retrospective study conducted in a university setting (Saveetha dental college and hospitals, Chennai). Thus the data available is of patients from the same geographic location and have similar ethnicity. Approval was obtained from the institutional Ethical committee (IEC). The ethical approval number for the present study is SDC/SIHEC/2020/DIASDATA/0619-0320.

**Sampling:** In this retrospective study, about 86,000 records of patients who reported to Saveetha Dental College from June 2019 to March 2020 were reviewed. The study population included 50 patients who had oral cancer with various forms of habits. Two examiners were involved in the study. Cross verification of data were done with the help of photographs. The internal validity of the study was carried out by analysing the age and gender of patients. The external validity was carried out by analysing the forms of habits.

**Data collection and tabulation:** Clinical examination, dental status and photographs of oral cancer patients with various forms of habits were assessed. Data related to age, gender, various forms of habits in oral cancer patients was collected. The study involved age groups divided into less than 50 years and greater than the 50 years. Incomplete or censored data was excluded from the study. Cross verification of data were done with the help of photographs. Two examiners were involved in the study. Data was entered in a methodical manner. Data was recorded and tabulated on Excel.

**Statistical Analysis:** After tabulation using MS Excel, the data was exported to IBM SPSS software [Version 19: IBM Corporation NY USA] for statistical analysis. Independent variables included age, gender. Dependent variables included various forms of habits. The statistical analysis was done by Pearson chi square test. Association of age, gender, various forms of habits in oral cancer patients were performed. Pearson chi square test was done to statistically analyze the data and was used to identify any significant level of association which was set at 0.05.

## RESULTS AND DISCUSSION

Out of the total 50 patients involved in the study it was observed that the highest number of oral cancer patients had Pan chewing as the most common form of habit (n=11;22%) followed by Pan chewing and Gutkha (n=4;8%), Gutkha (n=3;6%) Smoking (n=2;4%), and for about 30 patient the cause was unknown (n=30;60%) [Figure 1]. It was

also observed that it was predominant in above 50 years of age group (n=29;58%) compared to females (n=21;42%) Pearson's Chi square value =5.136; df =4 ;p-value: 0.274 (p>0.05); hence statistically not significant. [Figure 2] Oral cancer Patients with various forms of habits were more predominant in males (n=39;78%) when compared to females (n=11;22%) Pearson's Chi square value = 5.852; df =4 ;p-value: 0.210 (p>0.05); hence statistically not significant. Chi square test was done and association between age group, gender and various forms of habits was found to be statistically not significant (P>0.05). [Figure 3]

Thus, from this study it was observed that oral cancer incidence in patients based on forms of habits was predominant in above 50 years of age group with a high predominance in the males and Pan chewing being the most common form of habit. Further studies with a larger sample size are required to compare the efficacy of oral cancer based on various forms of habits.

Oral cancer is also the most common type of head and neck cancer in Taiwan, with the incidence rate among men being 10.9 times higher than that among women. By comparing the incidence rates of various cancers in Taiwan, it can be observed that oral cancer ranks fifth among the top 10 causes of deaths due to cancer and has been the fourth most common cancer among men for 12 consecutive years since 2003. Therefore, the prevention and control of oral cancer are regarded as key health issues in Taiwan<sup>41</sup>

Sex, age, educational level, marital status, presence or absence of diabetes, presence or absence of other cancers, comorbidity severity, smoking and betel nut chewing habits, and monthly salary influenced oral cancer incidence among individuals who had undergone oral cancer screening. Among the individuals screened, men had a higher risk of oral cancer incidence (hazard ratio (HR) =5.72)<sup>42</sup>. Subapriya *et al.* revealed in their study that combination of smoking, chewing, and alcohol had a 11.34-fold higher risk than independent risk of disease. Similar to earlier reports, the present retrospective study showed a total of 98.5% prevalence of buccal mucosa carcinoma with regard to risk habits of tobacco smoking, chewing, pan/areca nut chewing, and alcoholism. However, 39.4% had multihabits, which proved to be more aggressive than single habits. Further, compared with all habits, alcohol prevalence reported with lowest of 1% buccal mucosa carcinoma patients. This baseline characteristic on buccal cancer provides etiological clues for future hospital-/population-based epidemiological studies<sup>43</sup>.

Tobacco use and alcohol are known risk factors for cancers of the oral cavity. Estimates indicate 57% of all men and 11% of women between 15–49 years of age use some form of tobacco. Besides smoking, use of smokeless tobacco is also widely prevalent as noted in the use of Betel quid. This is very common and is accepted socially and culturally in many parts of India. Additionally, gutka, zarda, kharra, mawa, and khainni are all dry mixtures of lime, areca nut flakes, and powdered tobacco custom mixed by vendors<sup>44,45</sup>.

Among individuals with habit, 135 (42.18%) were tobacco chewers and smoking was prevalent in about 48 (15%). The prevalence was more in males 37 (15.22%). Amongst both males (44.03%) and females (48.05%) gingivo-buccal sulcus was the most common site. The least common site was seen to be lip with only 10 (3.12%) patients. Well differentiated squamous cell carcinoma was common in both males and females with 155 (63.78%) and 47 (61.03%) respectively. Poorly differentiated squamous cell carcinoma was only reported in 4 (1.64%) males<sup>46</sup>.

Early detection of these oral cancers will enable appropriate clinical management and monitoring. Moreover, improving the incidence, mortality, and survival rates of oral cancer requires a multi-tier structural approach that targets society, dentists, communities, and the individual. Our institution is passionate about high quality evidence based research and has excelled in various fields<sup>47-53</sup>. We hope this study adds to this rich legacy.

## CONCLUSION


Within the limitations of our study it was observed that oral cancer incidence in patients based on forms of habits was predominant in the above 50 years age group with a higher predilection for males and Pan chewing being the most common form of habit. Further studies with a larger sample size are required to compare the efficacy of oral cancer based on various forms of habits.

## REFERENCES


- i. Gundamaraju K, Kantheti LP, Naga SNV, et al. Prevalence of oral potentially malignant and malignant lesions at a tertiary level hospital in Hyderabad, India. *Journal of Dr. NTR University of Health Sciences* 2014; 3: 13.
- ii. Iodice S, Gandini S, Maisonneuve P, et al. Tobacco and the risk of pancreatic cancer: a review and meta-analysis. *Langenbeck's Archives of Surgery* 2008; 393: 535–545.
- iii. Reibel J. Tobacco and Oral Diseases. *Medical Principles and Practice* 2003; 12: 22–32.
- iv. Gandini S, Botteri E, Iodice S, et al. Tobacco smoking and cancer: A meta-analysis. *International Journal of Cancer* 2008; 122: 155–164.
- v. Radoi L, Luce D. A review of risk factors for oral cavity cancer: the importance of a standardized case definition. *Community Dentistry and Oral Epidemiology* 2013; 41: 97–109.
- vi. Warnakulasuriya S, Greenspan JS. *Textbook of Oral Cancer: Prevention, Diagnosis and Management*. Springer Nature, 2020.
- vii. International Agency for Research on Cancer. *Who Classification of Tumours of the Lung, Pleura, Thymus and Heart*. World Health Organization, 2015.
- viii. Mishra G, Pimple S, Shastri S. An overview of the tobacco problem in India. *Indian Journal of Medical and Paediatric Oncology* 2012; 33: 139.
- ix. Bhardwaj N, Daniel M, Srinivasan S, et al. Demographics, habits, and clinical presentation of oral cancer in Puducherry's population: An institutional experience. *Journal of Indian Academy of Dental Specialist Researchers* 2015; 2: 64.
- x. Rai HC, Ahmed J. Clinicopathological Correlation Study of Oral Squamous Cell Carcinoma in a Local Indian Population. *Asian Pacific Journal of Cancer Prevention* 2016; 17: 1251–1254.
- xi. Chaturvedi P, Chocolatewala N. Role of human papilloma virus in the oral carcinogenesis: An Indian perspective. *Journal of Cancer Research and Therapeutics* 2009; 5: 71.
- xii. Jikko A, Murakami S, Fujishita M, et al. Correlation between radiographic and histopathological findings in squamous cell carcinomas of the gingiva. *Oral Radiology* 1989; 5: 33–40.
- xiii. Murarkar SK, Gothankar JS, Doke P, et al. Manuscript title: Prevalence and determinants of undernutrition among under-five children residing in urban slums and rural area, Maharashtra, India: a community-based cross-sectional study. DOI: 10.21203/rs.3.rs-15651/v2.
- xiv. Agrawal M, Pandey S, Jain S, et al. Oral Cancer Awareness of the General Public in Gorakhpur City, India. *Asian Pacific Journal of Cancer Prevention* 2012; 13: 5195–5199.
- xv. Abhinav R, Selvarasu K, Maheswari G, et al. The patterns and etiology of maxillofacial trauma in South India. *Annals of Maxillofacial Surgery* 2019; 9: 114.
- xvi. Christabel A, Anantanarayanan P, Subash P, et al. Comparison of pterygomaxillary dysjunction with tuberosity separation in isolated Le Fort I osteotomies: a prospective, multi-centre, triple-blind, randomized controlled trial. *International Journal of Oral and Maxillofacial Surgery* 2016; 45: 180–185.
- xvii. Jesudasan JS, Abdul Wahab PU, Muthu Sekhar MR. Effectiveness of 0.2% chlorhexidine gel and a eugenol-based paste on postoperative alveolar osteitis in patients having third molars extracted: a randomised controlled clinical trial. *British Journal of Oral and Maxillofacial Surgery* 2015; 53: 826–830.
- xviii. Kumar S. KNOWLEDGE, ATTITUDE AND AWARENESS OF DENTAL UNDERGRADUATE STUDENTS REGARDING HIV/AIDS PATIENTS. *Asian Journal of Pharmaceutical and Clinical Research* 2017; 10: 175.
- xix. Kumar S. RELATIONSHIP BETWEEN DENTAL ANXIETY AND PAIN EXPERIENCE DURING DENTAL EXTRACTIONS. *Asian Journal of Pharmaceutical and Clinical Research* 2017; 10: 458.
- xx. Kumar S. THE EMERGING ROLE OF BOTULINUM TOXIN IN THE TREATMENT OF OROFACIAL DISORDERS: LITERATURE UPDATE. *Asian Journal of Pharmaceutical and Clinical Research* 2017; 10: 21.
- xxi. Kumar S, Rahman R. KNOWLEDGE, AWARENESS, AND PRACTICES REGARDING BIOMEDICAL WASTE MANAGEMENT AMONG UNDERGRADUATE DENTAL STUDENTS. *Asian Journal of Pharmaceutical and Clinical Research* 2017; 10: 341.
- xxii. Kumar S, Sneha S. KNOWLEDGE AND AWARENESS REGARDING ANTIBIOTIC PROPHYLAXIS FOR INFECTIVE ENDOCARDITIS AMONG UNDERGRADUATE DENTAL STUDENTS. *Asian Journal of Pharmaceutical and Clinical Research* 2016; 154.
- xxiii. Marimuthu M, Andiappan M, Wahab A, et al. Canonical Wnt pathway gene expression and their clinical correlation in oral squamous cell carcinoma. *Indian Journal of Dental Research* 2018; 29: 291.
- xxiv. Packiri S. Management of Paediatric Oral Ranula: A Systematic Review. *JOURNAL OF CLINICAL AND DIAGNOSTIC RESEARCH*. Epub ahead of print 2017. DOI: 10.7860/jcdr/2017/28498.10622.
- xxv. Patil SB, Durairaj D, Suresh Kumar G, et al. Comparison of Extended Nasolabial Flap Versus Buccal Fat Pad Graft in the Surgical Management of Oral Submucous Fibrosis: A Prospective Pilot Study. *Journal of Maxillofacial and Oral Surgery* 2017; 16: 312–321.
- xxvi. Patturaja K, Pradeep D. Awareness of Basic Dental Procedure among General Population. *Research Journal of Pharmacy and Technology* 2016; 9: 1349.
- xxvii. Rao TD, Santhosh Kumar MP. Analgesic Efficacy of Paracetamol Vs Ketorolac after Dental Extractions. *Research Journal of Pharmacy and Technology* 2018;

- 11: 3375.
- xxviii. Abhinav RP, Sweta VR, Ramesh A. Role of virtual reality in pain perception of patients following the administration of local anesthesia. *Annals of Maxillofacial Surgery* 2019; 9: 110.
- xxix. Jain SV, Vijayakumar Jain S, Muthusekhar MR, et al. Evaluation of Three-Dimensional Changes in Pharyngeal Airway Following Isolated Lefort One Osteotomy for the Correction of Vertical Maxillary Excess: A Prospective Study. *Journal of Maxillofacial and Oral Surgery* 2019; 18: 139–146.
- xxx. Neal RD, Allgar VL. Sociodemographic factors and delays in the diagnosis of six cancers: analysis of data from the 'National Survey of NHS Patients: Cancer'. *British Journal of Cancer* 2005; 92: 1971–1975.
- xxxi. Krishnan S, Pandian S, Kumar S A. Effect of bisphosphonates on orthodontic tooth movement—an update. *J Clin Diagn Res* 2015; 9: ZE01–5.
- xxxii. Ramesh Kumar KR, Shanta Sundari KK, Venkatesan A, et al. Depth of resin penetration into enamel with 3 types of enamel conditioning methods: a confocal microscopic study. *Am J Orthod Dentofacial Orthop* 2011; 140: 479–485.
- xxxiii. Felicita AS. Orthodontic management of a dilacerated central incisor and partially impacted canine with unilateral extraction - A case report. *Saudi Dent J* 2017; 29: 185–193.
- xxxiv. Kumar S. The emerging role of botulinum toxin in the treatment of orofacial disorders: Literature update. *Asian J Pharm Clin Res* 2017; 10: 21.
- xxxv. Felicita AS. Quantification of intrusive/retraction force and moment generated during en-masse retraction of maxillary anterior teeth using mini-implants: A conceptual approach. *Dental Press J Orthod* 2017; 22: 47–55.
- xxxvi. Sivamurthy G, Sundari S. Stress distribution patterns at mini-implant site during retraction and intrusion—a three-dimensional finite element study. *Prog Orthod* 2016; 17: 1–11.
- xxxvii. Sekar D, Lakshmanan G, Mani P, et al. Methylation-dependent circulating microRNA 510 in preeclampsia patients. *Hypertens Res* 2019; 42: 1647–1648.
- xxxviii. Johnson J, Lakshmanan G, M B, et al. Computational identification of MiRNA-7110 from pulmonary arterial hypertension (PAH) ESTs: a new microRNA that links diabetes and PAH. *Hypertens Res* 2020; 43: 360–362.
- xxxix. Jain RK, Kumar SP, Manjula WS. Comparison of intrusion effects on maxillary incisors among mini implant anchorage, j-hook headgear and utility arch. *J Clin Diagn Res* 2014; 8: ZC21–4.
- xl. Keerthana B, Thenmozhi MS. Occurrence of foramen of huschke and its clinical significance. *Research Journal of Pharmacy and Technology* 2016; 9: 1835–1836.
- xli. Hung L-C, Kung P-T, Lung C-H, et al. Assessment of the Risk of Oral Cancer Incidence in A High-Risk Population and Establishment of A Predictive Model for Oral Cancer Incidence Using A Population-Based Cohort in Taiwan. *International Journal of Environmental Research and Public Health* 2020; 17: 665.
- Shafique K, Mirza S, Vart P, et al. Areca nut chewing and systemic inflammation: evidence of a common pathway for systemic diseases. *Journal of Inflammation* 2012; 9: 22.
- Subapriya R, Thangavelu A, Mathavan B, et al. Assessment of risk factors for oral squamous cell carcinoma in Chidambaram, Southern India: a case-control study. *European Journal of Cancer Prevention* 2007; 16: 251–256.
- Gupta P, Pindborg J, Bhonsle RB, et al. INTERVENTION STUDY FOR PRIMARY PREVENTION OF ORAL CANCER AMONG 36 000 INDIAN TOBACCO USERS. *The Lancet* 1986; 327: 1235–1239.
- Malaowalla AM, Silverman S, Mani NJ, et al. Oral cancer in 57,518 industrial workers of Gujarat, India. A prevalence and followup study. *Cancer* 1976; 37: 1882–1886.
- Raj A, Ramesh G, Pathak S. Assessment of hospital based prevalence of oral cancer among population of Kanpur, Uttar Pradesh. *J Exp Ther Oncol* 2018; 12: 193–199.
- Vijayashree Priyadharsini J. In silico validation of the non-antibiotic drugs acetaminophen and ibuprofen as antibacterial agents against red complex pathogens. *J Periodontol* 2019; 90: 1441–1448.
- Ezhilarasan D, Apoorva VS, Ashok Vardhan N. Syzygium cumini extract induced reactive oxygen species-mediated apoptosis in human oral squamous carcinoma cells. *J Oral Pathol Med* 2019; 48: 115–121.
- Ramesh A, Varghese S, Jayakumar ND, et al. Comparative estimation of sulfiredoxin levels between chronic periodontitis and healthy patients - A case-control study. *J Periodontol* 2018; 89: 1241–1248.
- Mathew MG, Samuel SR, Soni AJ, et al. Evaluation of adhesion of Streptococcus mutans, plaque accumulation on zirconia and stainless steel crowns, and surrounding gingival inflammation in primary .... *Clin Oral Investig*, <https://link.springer.com/article/10.1007/s00784-020-03204-9> (2020).
- Sridharan G, Ramani P, Patankar S, et al. Evaluation of salivary metabolomics in oral leukoplakia and oral squamous cell carcinoma. *J Oral Pathol Med* 2019; 48: 299–306.
- Pc J, Marimuthu T, Devadoss P. Prevalence and measurement of anterior loop of the mandibular canal using CBCT: A cross sectional study. *Clin Implant Dent Relat Res*, <https://europepmc.org/article/med/29624863> (2018).
- Ramadurai N, Gurunathan D, Samuel AV, et al. Effectiveness of 2% Articaine as an anesthetic agent in children: randomized controlled trial. *Clin Oral Investig* 2019; 23: 3543–3550.






**Figure 1.** Bar graph depicts the frequency distribution of various forms of habit in patients with oral cancer. The X axis represents form of habit, Y axis represents the number of patients. The graph shows that the highest form of habit seen in oral cancer patients was Pan chewing.



**Figure 2.** Bar graph depicts the association between gender and number of oral cancer patients with various forms of habits. X axis represents gender, Y axis represents the number of oral cancer patients with various forms of habits. It was observed that majority of the patients form of habit was unknown. (Green) There was no association between gender and number of oral cancer patients with various forms of habits. Pearson's Chi square value = 5.136; df = 4 ; p-value: 0.274 (p > 0.05); hence statistically not significant.



**Figure 3:** Bar graph depicts the association between age and number of oral cancer patients with various forms of habits. X axis represents age, Y axis represents the number of oral cancer patients with various forms of habits. It was observed that majority of the patients form of habit was unknown. (Green) There was no association between age and number of oral cancer patients with various forms of habits. Pearson's Chi square value = 5.852; df = 4 ; p-value: 0.210 (p > 0.05); hence statistically not significant.