

attractive because, their ability to leave by products into atoms efficient and environmentally friendly, namely H₂O [6]. Then the reaction is generally followed by a carbocation addition to the double bond and subsequent protonated [7]. The use of FeCl₃ has shown the catalytic activity of alcohols against various nucleophiles [8]. Alkyne used in the formation of an alkenyl halide is 17 α -ethinylestradiol, where the compound is a drug commonly used in disease problems infertility, hormone hoarding, and osteoporosis [9]. Previous studies have shown that 17 α -ethinylestradiol (EE2) is highly resistant to oxidation in the environment due to the presence of an ethinyl group at position 17, and because it has a high C \equiv C bond energy, 17 α -ethinylestradiol (EE2) has an estrogenic ability 11-30 times stronger than estrogen (E2) [10]. Previous research involved alkynes and alcohols as the main reactants and HX as a source of halides and using the C-C coupling reaction. A compound formed from an addition of cyclohexanol at 17 α -ethinylestradiol with anhydrous ferric chloride (FeCl₃) salt medium is an estradiol derivative compound with chloride alkenyl group is at atom C-17.

2. RESEARCH METHODS

Material. 17 α -ethinylestradiol (Sigma aldrich) 98% purity, cyclohexane, anhydrous iron (III) chloride, dichloromethane pa, n-Hexane, chloroform pa, ethinyl acetate, aquadest, methanol, H₂SO₄, TLC plate F₂₅₄, silica gel 60 (70- 230 mesh), aluminum foil, cotton.

The equipment used was Beaker, spatula, measuring pipette, eppendorf micro pipette, dropper, stative, clamp, measuring cup, watch glass, capillary tube, vial, tweezers, spray bottle, TLC elution chamber, gravity chromatography column, analytical balance, fume hood, oven mert, rotary shaker, centrifuge tube.

Cross-Coupling Reaction

Anhydrous iron (III) chloride (16 mg, 0.1

mmol), cyclohexane (221 mg, 1.2 mmol), and dichloromethane solvent (3 mL) were mixed in a centrifuge tube by shaking. After being shaken for several minutes, the mixture was then added with 17 α -ethinylestradiol (132 mg, 1 mmol) and continued shaking for 48 hours. During the process, the reaction was monitored by testing thin layer chromatography.

Separation of compound

After the reaction was completed, the reaction mixture was separated by gravity column chromatography using n-hexane:ethinyl acetate as mobile phase with several ratios of 10:0; 9:1; 8.5:1.5; 8:2; 7:3.

Identification of Sample Fraction

The fraction result from the separation was further identified using the 1H-NMR (Nuclear Magnetic Resonance) testing technique.

3. RESULT AND DISCUSSION

Cyclohexanol Addition Reaction Stage in EE2 with FeCl₃.

The results of the reaction were carried out in the first step, namely the process of mixing the reaction between FeCl₃, cyclohexanol, and dichloromethane into a centrifuge tube and shaken using a Rotary Shaker for 30 minutes. This conditioning process is carried out with the aim of forming a FeCl₃-cyclohexanol complex, which causes the formation of a dimeric ether compound. After that, EE2 is added and can then immediately attack the dimeric FeCl₃-ether complex, where the triple bond acts as nucleophilic. At this stage, monitoring was carried out using thin layer chromatography (TLC). TLC analysis is an analysis of chemical components using the principle of absorbent and partition, where the chemical components will move following the mobile phase this is because the absorbent absorption of chemical components is not the same so that these components will move with different

distances [11]. The reaction was stopped when it was seen that there were other components separated from the standard

(there were other stains with different Rf from the standard) the Rf value of the TLC test results can be seen in Table 1.

Table 1. Rf value of TLC results from the reaction mixture

	Rf		
	EE2	S	X
CHCl ₃ (9.5) : EtOAc (0.5)	0.3	0.5	0.5
CHCl ₃ (9) : n-hexane (1)	0.3	0.31	0.31
n-hexane (7): EtOAc (3)	0.57	0.67	0.67

Description: S = Spike (Standard + sample)

X = sample

Separation of the Addition Reaction Mixture

Gravity column chromatography technique was performed with a column length of 30 cm and a diameter of 1.5 cm. ± 15 g of silica gel was used, then soaked for 1x24 hours using 100 ml of n-hexane as solvent. The eluent used for steroid and tripenoid compounds consisted of n-hexane and ethyl acetate in a ratio of 16:4; 17:3; and 18:2, where the n-hexane solution is a non-polar solvent and ethyl acetate is a semi-polar solvent [12]. In this research, TLC analysis was carried out on several solvents, then the eluent for column

chromatography was obtained, namely n-Hexane:Ethylacetate with several ratios, namely 20:0; 36:4; 34:6; 32:8; 14:6. The sample will be eluted based on the interaction with the stationary phase and the flow reaction rate with a time range of 6-14 seconds/drop, then it is accommodated into 60 vials and each vial holds 2 ml/vial. Furthermore, the eluate was grouped using hexane/ethyl acetate eluent (8:2). The Rf value of the TLC analysis on the separation of the reaction mixture before grouping can be seen in Table 2.

Table 2. Rf value of TLC analysis on eluate at intervals of 3 vials using n-hexane (8) : ethyl acetate (2) eluent.

No. Vial	Rf value	No. Vial	Rf value
03	.	36	0.5
06	0.94	39	-
09	0.94	42	0.22
12	0.94	45	0.22
18	.	48	0.22
21	.	51	0.22
24	0.5	54	0.22
27	0.44	57	0.22
30	0.52	60	-
33	0.48	EE2	0.22

From the existing Rf data, 3 groups of fractions were obtained, namely I (1-22); II (23-36); III (37-60) and fraction II were carried out in a further step, 1H-NMR test because fraction II showed a different stain appearance with the characteristics of the standard stain (EE2) and had a higher Rf value than the standard Rf.

Structure Determination

Fraction II was analyzed using a Nuclear Magnetic Resonance Proton (1H-NMR) instrument to determine the structure of the compound, using Chloroform-D (CDCl₃) solvent and measured at a frequency of 500 MHz. Table 3 shows the spectroscopic data for 1H-NMR fraction II by comparing the standard EE2 1H-NMR data and the comparison journal.

Table 3. Data tabulation of 1H-NMR fraction II with standards and comparison journals.

Proton position	EE2* δ H	Standard EE2 δ H	Fraction I δ H
1	7,05	7,088	7,517
2	6,50	6,569	7,509
4	6,43	7,945	7,450
6 α	2,7 ^a	2,722	2,349
6 β	2,7 ^b	2,718	2,335
7 α	1,25	1,280	1,269
7 β	1,76	1,718	1,670
8	1,30	1,375	1,269
9	2,04	2,025	2,000
11 α	2,28	2,299	2,281
11 β	1,30	1,389	1,319
12 α	1,76	1,712	1,605
12 β	1,65	1,692	1,670
14	1,31	1,399	1,239
15 α	1,65	1,692	1,670
15 β	1,31	1,399	1,239
16 α	2,10	2,182	2,272
16 β	1,86	1,858	1,319
18	0,75	0,857 ^s	0.814
		3,297 ^d	
20	3,31	(J = 5 Hz)	5,350
OH	5,29	4.334 ^s	4,137
			4,116

Exp*= comparison journals.

In steroids, ring alternation plays an important role. Chemical shift H 1.57 ppm; H 1.66 ppm ; H 2.33 ppm is predicted to be a proton signal contained in the cyclohexanol group [14]. The following interpretation of the chemical shift tabulation data in fraction II is as follows:

- The chemical shift of H 4.116 ppm indicates the presence of –OH group attached to the C-17 group.
- The chemical shift of H 4.137 ppm indicates the presence of an –OH group attached to the C-3 group.
- The aromatic proton signal appears at the H shift of 7.517; 7.509; 7,450 ppm
- The signal of the proton bound to the C-8 methyl is in the H shift of 0.814
- A shift of 5.350 identifies the presence of a proton attached to a double bond

(C-20).

The shift for the alkenyl group shifted in the range of 4.4-7.5 ppm [15]. The signal for C-20 in fraction II shows a shift at H 5,350. This indicates that there are no more alkyne

groups at C-20. From the tabulated data, a new compound derived from estradiol was obtained and the structure and name of the compound were obtained as shown in Figure 2.




Figure 2. 17 α -(19-chlorovinilic-20 cyclohexane)estradiol.

Addition Reaction Mechanism

FeCl₃ here acts as a strong Lewis acid and with this property, the oxygen atom is partially positively charged due to the attraction of the electron pair present in Iron(III). The oxygen atom in cyclohexanol is unstable because it binds to FeCl₃, then the oxygen atom affects the carbon-oxygen bond in the cyclohexyl ring to stabilize its partial charge. The proton that binds to oxygen is then released by means of the released hydroxyl anion which binds it to form a water molecule (H₂O) and a dimeric ether compound. In the presence of a

nucleophile, the ether is re-polarized by FeCl₃, resulting in a new benzylic carbocation and an iron (III) atom in FeCl₃ which has a partial negative charge then EE2 is added which in this case is nucleophilic. The ethinyl group which has an electron pair attacks the cyclohexyl carbon to form a vinylic carbocation and an iron complex. This vinyl carbocation then stabilizes itself by binding to one of the chlorines present in FeCl₃. The mechanism of the addition reaction is shown in

figure

4




Figure 4. Proposed Reaction Mechanism.

4. CONCLUSION

1. The addition reaction between 17 α -ethinylestradiol and cyclohexanol with FeCl₃ media produces a 17 α (19-chlorovinilic-20 cyclohexane)estradiol, where the alkyne group is converted to an alkenyl halide group. This is evidenced by a chemical shift in the proton C-20 which is 5.350 ppm.
2. The mechanism of formation of alkenyl halides by addition reaction begins with the formation of dimeric ether from cyclohexanol using FeCl₃ as a catalyst and produces an environmentally friendly by-product in the form of water (H₂O).

REFERENCES

1. Hotma R. Simbolon, R. J. (2021). Kajian Regiokimia Markovnikov Melalui Mekanisme Reaksi Hidrasi Alkuna Terminal Pada Etinilestradiol dengan Katalis FeCl₃. *FULLERENE Journal of Chemistry*. In Press
2. Biswas, S. D. (2009). New And Efficient iron Halide Mediated Synthesis Of Alkenyl Halides Through Coupling Of Alkynes And Alcohols. *European Journal Of Organic Chemistry*, 2354-2359.
3. Francis A. Carey, R. M. (2011). *Organic Chemistry Eighth Edition*. Americas, New York.: McGraw-Hill.
4. Brown, F. . (2011). *Organic Chemistry Sixth Edition*. Usa.
5. Furstner, A. A. (2002). rion-Catalyzed Cross-Coupling Reactiions Of Alkyl-Grignard Reagents With Aryl Chlorides, Tosylates, And Triflates. *Angew. Chem. nt.Ed*, 609-612.
6. Liu, Z. Q. (2009). An Efficient And Mild iron-Mediated Synthesis Of Alkenyl Halides Via Direct C-C Bond Formatiion Of Benzyl Alcohol And Aryl Alkynes. *Tetrahedron Letters Elsevier*, 1240-1242.
7. Fenga, W. D. (2018). iron-Catalyzed Olefin Synthesis By Direct Coupling Of Alkenes With Alcohols: A Dft investigation. *ComputatiionalAndTheoretical Chemistry 1143. Elsevier*, 36-42.
8. Jana, U. D. (2008). iron(III)-Catalyzed Addition Of Benzylic Alcohols To Aryl Alkynes- A New Synthesis Of Substituted Aryl Ketiones. *European Journal Of Organic Chemistry*, 5789-5804.
9. Wang, Y. D. (2014). Adsorptiion Behaviors Of 17 α -Ethinylestradiol n Sediment-Water System n Northern Taihu Lake, China. *Hindawi Publishing Corporatiion The Scieintific World Journal*, 1-6.
10. Szabo, L. D. (2020). Investigation Of The Sorption Of 17 α -Ethinylestradiol (EE2) on Soils Formed Under Aerobic And Anaerobic Conditions. *Elsevier*, 1-10.
11. Yohannes Alen, F. L. (2017). Analisis Kromatografi Lapis Tipis (KLT) dan Aktivitas Antihiperurisemia Ekstrak Rebung *Schizostachyum brachycladum Kurz* (Kurz) pada Mencit Putih Jantan. *Jurnal Sains Farmasi & Klinis*, 146-152.
12. Armeida Dwi Ridhowati Madjid, D. A. (2020). Variasi Komposisi Eluen pada Isolasi dan Triterpenoid Alga Merah *Eucheuma cottoinii* dengan Kromatografi Kolom Basah. *ALCHEMY: JOURNAL OF CHEMISTRY*, 35-40
13. José A.A. França1, A. N.-V. (2016). Complete Nmr Assignment And Cionformational Analysis Of 17-A-Ethinylestradiol By Using RdcS Obtained n Grafted Graphene Oxide.
14. Dayang Irma Suryaini, T. A. (2017). Karakteristik Senyawa Steroid Dari Fraksi Diklorometana Ranting Durian Klawing (*Durio graveolens Becc.*). *JKK*, 45-48
15. Webster, S. D. (1981). *Spectrometric identification Of Organic Compounds* .
16. Kolinug, R. Y. (2016). Reaksi Direct-Coupling 17 α -Etinilestradiol Dan Sikloheksanol Dengan Besi (III) Klorida Anhidrat Sebagai Media Reaksi .