



## PHYTOCHEMICAL ANALYSIS OF PROSOPIS JULIFLORA FOR ITS ANALGESIC PROPERTY

Dr. K.Hemamalini and <sup>2</sup>Dr. Sadanandam palle,

\* Professor and Principal, Department of Pharmacology, Swami Vivekananda Institute of Pharmaceutical Sciences, Vangapally (v), Yadagirigutta (M), Yadadiri, Bhongiri(D), 508286, <sup>2</sup> Professor, Chemistry department , Malla reddy Engineering college , Hyderabad

Correspondence to: [rkhemamalini@gmail.com](mailto:rkhemamalini@gmail.com), 9553601248

### Abstract :

**Background:** The most economical and ecological nature of *Prosopis juliflora* with its medicinal value in arid and semi-arid zones of the world for its different properties against inflammation, it is a tree, belongs to the family Leguminosae (Fabaceae).

**Aims and Objectives:** To evaluate the analgesic activity of methanolic extract of dried leaves of *Prosopis juliflora* Linn.

**Materials and Methods:** Adult male Wistar rats (100–150 g body weight) were used in this study. Methanolic extract of *Prosopis juliflora* Linn was used to evaluate acute analgesic activity by tail flick method by oral administration at doses of 300, and 500 mg/kg body weight in healthy albino rats.

**Result:** In acute studies, the methanolic extract showed significantly and dose-dependently reaction time but after 1 hour the reaction time was absent. Statistical analysis was carried out by one-way ANOVA, followed by Turkey's test.

**Conclusion:** methanolic extract of *Prosopis juliflora* Linn possesses analgesic activity in a dose-dependent manner in thermal induced model.

**KEY WORDS:** *Prosopis juliflora*; Analgesic; tail flick model.

### INTRODUCTION:

Environmental like natural plants always stands as golden mark for the outstanding phenomena of symbiosis. The medicinal property of the natural obtaining materials serves humans to treat the illness and to maintain their body in a healthy way since from ancient period. For the treatment of human diseases different products obtained from natural sources like plants, minerals and even from animal sources to<sup>1</sup>. For an individual and their communities to maintain a healthy life they need the god gifted natural products. The products obtained from the plant sources shows a therapeutic property to live a long life and this is practiced in many countries<sup>2</sup>. The plant product was used as a medicine since from the 3 or 4 generation since our ancient. Plant products are the triumph of popular therapeutic diversity<sup>3</sup>. In recent times after the pandemic the world is now moving towards the herbal medicine or system, which can then properly fight the invading foreign particles, and help to destroy offending pathogens with less toxic effects<sup>4</sup>. In the early 1970's WHO had encouraged government for effective usage of tribal

information from the elder tribes and their local knowledge of herbal medicines for disease prevention and health promotion<sup>5</sup>. These information was documented for few plants<sup>6</sup> in different city libraries according to the region of plant availability. There are many species in the plant kingdom which contain medicinal value and these are to be discovered yet. All over the world the people are aware that India is rich source of herbal plants with medicinal property and yet the research is very important to evaluate it. In last two decades many people got interest towards the plant research and this was increased after the pandemic situation. From the academic view point it is apparent that students of botany, phytochemistry and pharmacology have now also come to expect some in –depth studies relating to medicinal plants. The new plant product or the drug obtained from the plant is brought to the market after several procedures where the new modern techniques and different pharmacological testing procedure help for it. For these new testing procedure the pharmacist play a vital role to evaluate the macroscopically and histological characters of the dried plant, which is tried through the chromatographic and other procedures for the identification and determination of purity of the preparation supplied. The plants used in the traditional style medicine of India and China as now receiving much scientific attention<sup>7</sup>.

An analgesic, or painkiller, is group of drugs used to relief from pain<sup>8</sup>. Analgesic drugs act through peripheral and central nervous systems. The examples are non-steroidal anti-inflammatory drugs [NSAIDs] such as the salicylate, and opioid drugs such as morphine and opium. An analgesic is a drug that selectively relieves pain by acting in the CNS or on peripheral pain mechanisms, without significantly altering consciousness. Pain is a warning signal, primarily protective in nature, but causes discomfort and suffering; may even be unbearable and incapacitating. Excessive pain may produce other effects- sinking sensation, apprehension, sweating, nausea, palpitation, rise or fall in BP, tachypnoea. Analgesics relieve pain as a symptom, without affecting its cause<sup>9</sup>.

#### **MATERIALS AND METHODS:**

**PLANT MATERIALS:** *Prosopis juliflora* belongs to the family Leguminosae (Fabaceae), sub-family Mimosoideae, with 44species out of which 40 are native to the Americas, three to Asia and one to Africa. The tropical Andean region is home to six species and eight species are found in the texas area, seven of them being endemic. These species have several properties such as soil binders, sand stabilizers, as well as its ability to grow in the poor soils. It is a shrub or tree which is 8-12 meters long and Grow to a height of 12 meters (39 ft). *P. juliflora* has trunk diameter of up to 1.2 meters (3.9 ft).

Its leaves are deciduous, geminate-pinnate, light green, with 12 to 20 leaflets. The Leaves appear short after leaf development. The tree reproduces solely by way of seeds, not vegetative. Seeds are spread by cattle and other animals, which consume the seed pods and spread the seeds in their droppings. The tree is said to have been introduced to Srilanka in the 19th century, where it is now known as vanni-andara, or katu andara in Sinhala. It is claimed that *P.juliflora* was recognized even as a holy tree in ancient India, but this is most likely confusion with *Prosopis*

cineraria<sup>10</sup>. In the western extent of its range in Ecuador and Peru, *Prosopis juliflora* readily hybridizes with *Prosopis pallida* and can be difficult to distinguish from this similar species or their interspecific hybrid strains<sup>11</sup>.

The various chemical agents that are present show the medicinal value that alters certain physiological actions in the human body. The several bioconstituents present in the plant are Terpenes, alkaloids, Flavonoids and phenolic compounds. Terpenes are used as insecticides and their pharmacological properties against some bacteria, fungal, earth worms, mosquitoes<sup>12</sup>. The Extracts of *P. juliflora* seeds and leaves have several in vitro pharmacological effects such as anti-bacterial, anti-fungal and anti-inflammatory properties<sup>13</sup>.

Since it is a main source as fuel for both urban and rural, this plant provides more than 90% of the fuel wood in some Indian villages, because *P. juliflora* wood has excellent burning qualities. Thus, it is called wooden anthracite. The wood obtained from this plant doesn't need storage and drying process<sup>14</sup>. *Prosopis juliflora* (Sw.) DC contains many alkaloids like juliflorine, julifloricine and julifloridine, juliprosine, juliprosinine and juliflorinine and show some responds the biological activity.

**Preparation of Plant Extract:** We have collected methanolic extract of *Prosopis juliflora* through Soxhlet apparatus by hot continuous extraction method. The use of commercially available Soxhlet apparatus is a convenient way to prepare crude plant extract. The dried and powdered drug was packed. Soxhlet apparatus works automatically and it is an continuous method that does not require further manipulation. It is not a time-consuming method and as for a standard-sized sample (50 g), extraction time is 48 h. The yield of the aqueous extract was 9.52%. The extract was stored in refrigerator until further studies.

**Drugs:** Ibuprofen, Diclofenac sodium, Aspirin (Cipla), acetic acid (ASES Chemical Works, Jodhpur), and Sodium chloride (ASES Chemical Works).

**Procurement of Animals:** Male Wistar rats weighing (100–150 g) were obtained. All animals were housed in ventilated cages and fed with a normal pellet diet and water ad libitum. All experiments were performed as per the ethical guidelines for investigations of experimental plant in conscious animal. Research protocol was approved by the Animal Ethics Committee.

#### **Anti-nociceptive Activity after Acute Administration by Tail-Flick Test**

Antinociceptive (analgesic) activity for the plant extract was evaluated by using tail-flick method. The rat where chosen for the study and divided into different groups in each group 6 animals were taken. Group I received the vehicle and served as control, group II served as a standard and group III received the plant extract. The animal rat was taken and the distal end of the rat was immersed in warm water maintained at 50°C. Only 5cm of the rat tail portion was taken and immersed in the water. The reaction time (in seconds) was the time taken by the rat to flick its tail due to pain. The first reading was omitted and reaction time was taken as the average of the next two readings. The reaction time was recorded before (0 min) and at 15, 30, 45, and 60 min after the administration of the treatments. The maximum reaction time was fixed at 15 sec

to prevent any tail tissue injury. If the reading exceeds 15 sec, it would be considered as maximum analgesia. The maximum possible analgesia (MPA) was calculated as follows:

MPA =  $\frac{\text{Reaction time for treatment} - \text{reaction time for saline}}{15 \text{ sec} - \text{reaction time for saline}} \times 100$ .

### Statistical Analysis:

The results are expressed as mean  $\pm$  SD (n = 6). Statistical significance was determined by ANOVA and subsequent Turkey's test. P values less than 0.05 were considered as indicative of significance.

### Results and Discussion:

The results of the analgesic activity of the methanol extract of the galls of *Prosopis juliflora* are shown in Table.1. Rats treated with normal saline (negative control) did not show any significant difference in the reaction time on tail-flick throughout the 60 min observation. In comparison with the baseline values within the same treatment groups, the increase in reaction time at different time points significantly differed ( $P < 0.05$ ) for morphine sulfate only. Duration of the reaction time in morphine sulfate and extract treated animals was significantly higher compared to saline treated animals, except for the extract group at 60 min. The highest reaction time for the extract treated group was 8.0 sec at 30 min, while it was 4.4 sec and 11.9 sec for saline and morphine sulfate groups, respectively. At all time points, the tail-flick latency time differed significantly between the extract and morphine sulfate groups, being greater for the latter group. No significant difference in reaction time was observed between the extract and sodium salicylate. Observation in rats treated with sodium salicylate did not give any significant analgesic effect in comparison with baseline values, saline, or extract (except for 30 min after treatment).

**TableNo:1 Analgesic effect of methanolic extract from the *Prosopis juliflora* by tail flick method in rats**

S.No	Treatments	Reaction time in seconds (mean $\pm$ SEM)				
		0 min	15 min	30 min	45 min	60 min
1	Control (normal saline)	4.25 $\pm$ 0.57	4.50 $\pm$ 0.34	4.42 $\pm$ 0.45	4.58 $\pm$ 0.44	5.17 $\pm$ 0.80
2	Morphine sulfate	6.50 $\pm$ 1.22	11.04 $\pm$ 0.73 <sup>*ab</sup>	10.92 $\pm$ 0.84 <sup>*ab</sup>	11.83 $\pm$ 0.35 <sup>*ab</sup>	13.33 $\pm$ 0.83 <sup>*ab</sup>
3	Sodium salicylate	4.13 $\pm$ 0.54	5.19 $\pm$ 0.57	6.75 $\pm$ 0.62 <sup>*a</sup>	5.65 $\pm$ 0.56	6.79 $\pm$ 1.24
4	MEPJ	6.57 $\pm$ 0.85 <sup>a</sup>	7.04 $\pm$ 0.67 <sup>a</sup>	8.04 $\pm$ 0.73 <sup>a</sup>	7.00 $\pm$ 0.92 <sup>a</sup>	6.57 $\pm$ 0.86

All values by Student's *t*-test, significant at  $P < 0.05$ , and SEM = standard error mean. <sup>\*</sup> $P < 0.05$  versus baseline of the respective treatment, <sup>a</sup>  $P < 0.05$  treatment versus control, <sup>b</sup>  $P < 0.05$  extract versus morphine sulfate, extract versus sodium salicylate was not significant at all time points.

Analgesics are drugs that act on peripheral or central nervous selectively relieve pain without significantly altering consciousness. Centrally acting analgesics act by raising the threshold for pain and also altering the physiological response to pain. On the other hand, peripherally acting analgesics act by inhibiting the generation of impulses at chemoreceptor site of pain. The animal models employed for screening of analgesic activity in this study are pain-state models using thermal stimuli which include tail-flick methods. Both methods are useful in illustrating centrally mediated antinociceptive responses which focus generally on changes above the spinal cord level. While the tail-flick method mediates a spinal reflex to a nociceptive stimulus, hot plate method involves higher brain functions and is regarded a supraspinally organized response.

In tail-flick model, the methanol extract from the *Prosopis juliflora* exhibited significant analgesic activity by increasing the reaction time of the rats compared to control (saline treated rats) at all time points, except at 60 min. Sodium salicylate and morphine sulfate were used as reference drugs, which are considered mild and moderate to severe analgesics, respectively. In comparison with control, morphine produced the most significant antinociception effect during all observation times, followed by the extract, while no significant analgesic effect was observed for sodium salicylate. The tail-flick method is based on the observation that morphine-like

compounds are selectively able to prolong the reaction time of typical tail-withdrawal effect in rats. This method is also useful in differentiating central opioid-like analgesics from peripheral analgesics. Analgesic drugs which are centrally acting elevate pain threshold of animals towards heat and pressure. Therefore, the analgesic effect of the extract on this pain-state model indicates that it might be centrally acting. With reference to the MPA value, the analgesic effects of both the extract and morphine sulfate were evident within 15 min following intraperitoneal administration. However, the extract showed short-lived analgesia as the MPA gradually decreased after 30 min compared to morphine sulfate. The tail-flick latency of the extract at all time points was less than that of reference drug, morphine sulfate, which is a slow onset opioid with long duration of action [17]. Although there was no significant analgesic effect between the reaction time of the extract and sodium salicylate, the extract exhibited a non-significant trend of higher reaction time compared to sodium salicylate. Both treatments produced comparable reaction times, suggesting that the galls of *Quercus infectoria* could be a better natural alternative for mild pain relief.

A number of alkaloids, flavanoids, steroids, and tannin isolated from medicinal plants have been reported to possess significant analgesic activity. The major constituent of the galls from *Prosopis juliflora* is tannin, which comprises up to 60% of its total content. Thus, analgesic activity observed with this extract might be attributed to the presence of this compound. Furthermore, there are reports on the role of tannin in analgesic activity. According to, preliminary phytochemicals which were screened from *Prosopis juliflora* L. including tannin might be responsible for the observed analgesic activity. Another research suggested that the presence of tannin and flavonoid in the methanol extract of *Prosopis juliflora* leaves seems to inhibit prostaglandin synthesis and exerts the anti-inflammatory and analgesic effects.

**Conclusion:** In this study, methanolic extract of MEPJ (500mg/kg, p. o.) significantly reduced the number of acetic acid-induced writhing and significantly increased the latency of paw licking in hot plate method.

**ACKNOWLEDGEMENT:** We like to thank the management of swami Vivekananda institute of pharmaceutical sciences who provided us the necessary facilities to make this research work in a successful way.

## REFERENCES

1. Chopra. R.N., Nayar. S.L., Chopra. I.C., "In Glossary of Indian medicinal plants", CSIR, New Delhi, 1<sup>st</sup> ed, 1956, 197.
2. The Ayurvedic Pharmacopoeia of India, "Ministry of health and family welfare Department and Indian syleavesof medicine and homeopathy", New Delhi, 11, (1), 1999, 137-140.
3. Yue- Zhong Shu., "Recent natural products based drug development: A Pharmaceutical Industry Perspective", **J. Nat. Prod.** 61, 1998, 1053-71.

4. Mukeshwar Pandey, Mousumi Debnath, Shobit Gupta, Surender K, Chikara, Phytomedicine: An Ancient approach turning into future potential source of therapeutics, **J, Pharmacog. Phytotherapy**, 3(3), 2011, 27-37.
5. Ravishankar. B., Shukla. V.J., "Indian syleavesof medicine: A brief profile, **African Journal Traditional complement alternative medicine**, 4(3), 2007, 319-337.
6. Kaido. T.L., Veale. D.J.H., Havlik. I., and Rama. D.B.K., **J. Ethnopharm.** 55, 1997, 185-191.
7. Trease. G.E., and Evans. W.C., **Pharmacognosy.**, 13<sup>th</sup> ed., 1992, 3-4.
8. Tambaro S, Reali R, Volonterio A, Zanda M, Olimpieri F, Pinna GA, Lazzari P [2013] NESS002 ie: A new fluorinated thiolendopeptidase inhibitor with antinociceptive activity in an animal model of persistent pain, **Pharmacology, Biochemistry and Behavior**, pp. 1-7.
9. Tripathi KD [2008]. Essentials of Medical Pharmacology, Sixth edition, Jaypee brothers medical publishers [p] ltd. pp.453.
10. H. Shiferaw Prosopis juliflora: the paradox of the dryland ecosystems, afar region, Ethiopia Wageningen, The Netherlands.
11. K.Saraswathi,NArunNagendran.Karthigaichamy R and Chandrasekaran S; Livelihood support from an invasive species Prosopisjuliflora.,**INT J CURR SCI**, 31-36 ,2012.
12. V. E. Vallejo, Z. Arbeli, W. Terán, N. Lorenz, R. P. Dick, Fabioroldan, Effect of land management and Prosopis juliflora (Sw.) DC trees on soil microbial community and enzymatic activities in intensive silvopastoral systems ofColombia. *Agriculture, Ecosystems and Environment* 150 (2012) 139– 148,Elsivier .
13. J Haji and A Mohammed Economic impact of Prosopis juliflora on agropastoralhouseholds of Dire Daw **Afri.J. of Agricul.Res**, 8(9), 768-779, 2013.
14. McKay GA, Reid JL, Walters MR. *Lecture Notes: Clinical Pharmacology and Therapeutics*. 8th edition. Blackwell Publishing; 2010.
15. Kilimozhi D, Parthasarathy V, Jayanth MNV, Manavalan R. Antinociceptive, antipyretic and anti-inflammatory effects of *Clerodendrum phlomidis* in mice and rats. *International Journal of Biological and Chemical Sciences*. 2009;3(3):504–512.